

FORM PTO-1390
(REV 10-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

15280-3511US

US APPLICATION NO. (if known, see 37 CFR 1.5)

097786666

INTERNATIONAL APPLICATION NO.

PCT/US99/20541

INTERNATIONAL FILING DATE

07 September 1999

PRIORITY DATE CLAIMED

08 September 1998

TITLE OF INVENTION PTERIDINE NUCLEOTIDE ANALOGS

APPLICANT(S) FOR DO/EO/US HAWKINS, Mary E., PFLEIDERER, Wolfgang,
BALIS, Frank


Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)).
4. ☒ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has **NOT** expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

International Preliminary Examination Report
International Search Report

U.S. APPLICATION NO. 09/786666		INTERNATIONAL APPLICATION NO. PCT/US99/20541		ATTORNEY'S DOCKET NUMBER 15280-3511PC	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1000.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	45 - 20 =	25	X \$18.00	\$ 450.00	
Independent claims	3 - 3 =	0	X \$80.00	\$	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$ 1310.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
SUBTOTAL =				\$ 1310.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$ 1310.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				\$	
TOTAL FEES ENCLOSED =				\$ 1310.00	
				Amount to be refunded:	\$
				charged:	\$
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed.					
b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>20-1430</u> in the amount of \$ <u>1310.00</u> to cover the above fees. A duplicate copy of this sheet is enclosed.					
c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. _____. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Joseph R. Snyder Townsend and Townsend and Crew LLP 2 Embarcadero Center, 8th Floor San Francisco, CA 94111-3834				 SIGNATURE: Kenneth A. Weber NAME 31,667 REGISTRATION NUMBER	



#5

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to:

Assistant Commissioner for Patents
Washington, D.C. 20231

On October 1, 2001

TOWNSEND and TOWNSEND and CREW LLP

By: Debra Ann DeBello
Debra Ann DeBello

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

HAWKINS et al.

Application No.: 09/786,666

Filed: March 7, 2001

For: PTERIDINE NUCLEOTIDE
ANALOGS

Examiner: Not yet assigned

Art Unit: Not yet assigned

RESPONSE UNDER 37 C.F.R. §§ 1.821-
1.825

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

This communication is being submitted in response to the Notification of Defective Response and Notice to Comply with Requirements for Patent Applications Containing Nucleotide Sequence and/or Amino Acid Sequence Disclosures, 37 C.F.R. §§ 1.821-1.825, mailed August 1, 2001. Applicants respectfully request reconsideration and further examination of the above-referenced patent application in view of the remarks presented herein.

REMARKS

The sequence listing was alleged to be in non-compliance with 37 CFR 1.821-1.825 for failing to comply with its requirements, and for the content of the computer readable form failing to comply with the requirements of 37 CFR 1.822 and/or 1.832. Applicant respectfully traverses. The Raw Sequence Error Report reports use of

09786666-061501

"n", where "n" must be equal to a nucleotide or a modified nucleotide. The Specification defines the fluorescent nucleotide analogs as the compounds listed where "n" was defined in the Sequence Listing. Thus, "n" is equal to a modified nucleotide.

To specifically detail the locations in the Specification where these definitions occur, reference should be made to:

- 1) Table 2 on page 39, where the sequences containing an "F" symbol occur, and the columns labeled "w/6MAP" and "w/DMAP" indicated which analogs are used.
- 2) The paragraph beginning on page 39, line 10, describing Table 2, where it states "in each row, F denotes the position of the probe within the sequence." Oligonucleotides containing 6MAP are number from 21 to 28 and those containing DMAP are numbered from 31 to 38.
- 3) Table 1 on page 38, following line 15, where "6MAP" and "DMAP" are defined as "compound 11" and "compound 12", respectively.
- 4) Example 4 A, page 33, lines 23-25, which gives the chemical name for compound 11, and Example 4B, page 34, lines 9-11, where the chemical name for compound 12 are described.


The Sequence Listing defines each "n", for the "modified base", in the "F" positions from each of the sequences found in Table 2 exactly as defined on pages 33 and 34. Therefore, Applicant respectfully requests that the "Notification of Defective Response" be withdrawn.

CONCLUSION

In view of the foregoing, Applicant believes the Sequence Listing now pending in this Application is proper. A first Action on the merits at an early date is respectfully requested.

If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 925-472-5000.

Respectfully submitted,



Joseph R. Snyder
Reg. No. 39,381

TOWNSEND and TOWNSEND and CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, California 94111-3834
Tel: (415) 576-0200
Fax: (415) 576-0300
SLW
SF 1274311 v1

09786666-001501



VW

RAW SEQUENCE LISTING ERROR REPORT

The Biotechnology Systems Branch of the Scientific and Technical Information Center (STIC) detected errors when processing the following computer readable form:

Application Serial Number: 09/786,666

Source: PCT 09

Date Processed by STIC: 7/5/2001

THE ATTACHED PRINTOUT EXPLAINS DETECTED ERRORS.

PLEASE FORWARD THIS INFORMATION TO THE APPLICANT BY EITHER:

- 1) INCLUDING A COPY OF THIS PRINTOUT IN YOUR NEXT COMMUNICATION TO THE APPLICANT, WITH A NOTICE TO COMPLY or,
- 2) TELEPHONING APPLICANT AND FAXING A COPY OF THIS PRINTOUT, WITH A NOTICE TO COMPLY

FOR CRF SUBMISSION QUESTIONS, PLEASE CONTACT MARK SPENCER, 703-308-4212.

FOR SEQUENCE RULES INTERPRETATION, PLEASE CONTACT ROBERT WAX, 703-308-4216.

PATENTIN 2.1 e-mail help: patin21help@uspto.gov or phone 703-306-4119 (R. Wax)

PATENTIN 3.0 e-mail help: patin3help@uspto.gov or phone 703-306-4119 (R. Wax)

TO REDUCE ERRORED SEQUENCE LISTINGS, PLEASE USE THE CHECKER VERSION 3.0 PROGRAM, ACCESSIBLE THROUGH THE U.S. PATENT AND TRADEMARK OFFICE WEBSITE. SEE BELOW:

Checker Version 3.0

The Checker Version 3.0 application is a state-of-the-art Windows based software program employing a logical and intuitive user-interface to check whether a sequence listing is in compliance with format and content rules. Checker Version 3.0 works for sequence listings generated for the original version of 37 CFR §§1.821 - 1.825 effective October 1, 1990 (old rules) and the revised version (new rules) effective July 1, 1998 as well as World Intellectual Property Organization (WIPO) Standard ST.25

Checker Version 3.0 replaces the previous DOS-based version of Checker, and is Y2K-compliant. Checker allows public users to check sequence listings in Computer Readable form (CRF) before submitting them to the United States Patent and Trademark Office (USPTO).

Use of Checker prior to filing the sequence listing is expected to result in fewer errored sequence listings, thus saving time and money.

Checker Version 3.0 can be down loaded from the USPTO website at the following address:

<http://www.uspto.gov/web/offices/pac/checker>

PCT

RAW SEQUENCE LISTING

DATE: 07/05/2001

PATENT APPLICATION: US/09/786,666

TIME: 15:44:04

Input Set : A:\Nih-3511.app

Output Set: N:\CRF3\07032001\I786666.raw

Does Not Comply
Corrected Diskette Needed

3 <110> APPLICANT: Hawkins, Mary E.
 4 Pfleiderer, Wolfgang
 5 Balis, Frank
 6 The Government of the United States of America
 7 as represented by The Secretary of the
 8 Department of Health and Human Services
 9
 10 <120> TITLE OF INVENTION: Pteridine Nucleotide Analogs
 11 <130> FILE REFERENCE: 015280-351100US
 12 <140> CURRENT APPLICATION NUMBER: US 09/786,666
 C--> 13 <141> CURRENT FILING DATE: 2001-06-15
 14
 15 <150> PRIOR APPLICATION NUMBER: US 60/099,487
 16 <151> PRIOR FILING DATE: 1998-09-08
 17 <150> PRIOR APPLICATION NUMBER: WO PCT/US99/20541
 18 <151> PRIOR FILING DATE: 1999-09-07
 19 <160> NUMBER OF SEQ ID NOS: 22
 20 <170> SOFTWARE: PatentIn Ver. 2.1
 21 <210> SEQ ID NO: 1
 22 <211> LENGTH: 21
 23 <212> TYPE: DNA
 24 <213> ORGANISM: Artificial Sequence
 25 <220> FEATURE:
 26 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
 27 PTR21
 28 <220> FEATURE:
 29 <221> NAME/KEY: modified_base
 30 <222> LOCATION: (7)
 31 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
 32 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
 33 N-diisopropyl)phosphoramidite (6MAP, compound 11)
 34
 35 <400> SEQUENCE: 1
 W--> 36 gtgtggnaaa tctctagcag t
 37 <210> SEQ ID NO: 2
 38 <211> LENGTH: 21
 39 <212> TYPE: DNA
 40 <213> ORGANISM: Artificial Sequence
 41 <220> FEATURE:
 42 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
 43 PTR22
 44 <220> FEATURE:
 45 <221> NAME/KEY: modified_base
 46 <222> LOCATION: (10)
 47 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
 48 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
 49 N-diisopropyl)phosphoramidite (6MAP, compound 11)
 50
 51 <400> SEQUENCE: 2
 W--> 52 gtgtgggaan tctctagcag t
 53 <210> SEQ ID NO: 3
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100

'n' must be equal to a
 nucleotide or a modified
 nucleotide

21

RAW SEQUENCE LISTING

PATENT APPLICATION: US/09/786,666

DATE: 07/05/2001

TIME: 15:44:04

Input Set : A:\Nih-3511.app

Output Set: N:\CRF3\07032001\I786666.raw

```

68 <211> LENGTH: 21
69 <212> TYPE: DNA
70 <213> ORGANISM: Artificial Sequence
72 <220> FEATURE:
73 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
74     PTR23
76 <220> FEATURE:
77 <221> NAME/KEY: modified_base
78 <222> LOCATION: (16)
79 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
80     beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
81     N-diisopropyl)phosphoramidite (6MAP, compound 11)
83 <400> SEQUENCE: 3
W--> 84 gtgtggaaaa tctcngcag t 21
87 <210> SEQ ID NO: 4
88 <211> LENGTH: 21
89 <212> TYPE: DNA
90 <213> ORGANISM: Artificial Sequence
92 <220> FEATURE:
93 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
94     PTR24
96 <220> FEATURE:
97 <221> NAME/KEY: modified_base
98 <222> LOCATION: (7)
99 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
100     beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
101     N-diisopropyl)phosphoramidite (6MAP, compound 11)
103 <400> SEQUENCE: 4
W--> 104 actgctngag attttccaca c 21
107 <210> SEQ ID NO: 5
108 <211> LENGTH: 21
109 <212> TYPE: DNA
110 <213> ORGANISM: Artificial Sequence
112 <220> FEATURE:
113 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
114     PTR25
116 <220> FEATURE:
117 <221> NAME/KEY: modified_base
118 <222> LOCATION: (9)
119 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
120     beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
121     N-diisopropyl)phosphoramidite (6MAP, compound 11)
123 <400> SEQUENCE: 5
W--> 124 actgctagng attttccaca c 21
127 <210> SEQ ID NO: 6
128 <211> LENGTH: 21
129 <212> TYPE: DNA
130 <213> ORGANISM: Artificial Sequence
132 <220> FEATURE:

```


RAW SEQUENCE LISTING
PATENT APPLICATION: US/09/786,666

DATE: 07/05/2001
TIME: 15:44:04

Input Set : A:\Nih-3511.app
Output Set: N:\CRF3\07032001\I786666.raw

133 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
134 PTR28
136 <220> FEATURE:
137 <221> NAME/KEY: modified_base
138 <222> LOCATION: (11)
139 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
140 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
141 N-diisopropyl)phosphoramidite (6MAP, compound 11)
143 <400> SEQUENCE: 6
W--> 144 actgctagcc nttttccaca c *See page 1* 21
147 <210> SEQ ID NO: 7
148 <211> LENGTH: 19
149 <212> TYPE: DNA
150 <213> ORGANISM: Artificial Sequence
152 <220> FEATURE:
153 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
154 HP21
156 <220> FEATURE:
157 <221> NAME/KEY: modified_base
158 <222> LOCATION: (10)
159 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
160 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
161 N-diisopropyl)phosphoramidite (6MAP, compound 11)
163 <400> SEQUENCE: 7
W--> 164 attccacaan gccgtgtca 19
167 <210> SEQ ID NO: 8
168 <211> LENGTH: 21
169 <212> TYPE: DNA
170 <213> ORGANISM: Artificial Sequence
172 <220> FEATURE:
173 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
174 HP22
176 <220> FEATURE:
177 <221> NAME/KEY: modified_base
178 <222> LOCATION: (11)
179 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
180 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
181 N-diisopropyl)phosphoramidite (6MAP, compound 11)
183 <400> SEQUENCE: 8
W--> 184 agaggtgtcc ncctgtggag a 21
187 <210> SEQ ID NO: 9
188 <211> LENGTH: 21
189 <212> TYPE: DNA
190 <213> ORGANISM: Artificial Sequence
192 <220> FEATURE:
193 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
194 HP23
196 <220> FEATURE:
197 <221> NAME/KEY: modified_base

RAW SEQUENCE LISTING
PATENT APPLICATION: US/09/786,666

DATE: 07/05/2001
TIME: 15:44:04

Input Set : A:\Nih-3511.app
Output Set: N:\CRF3\07032001\I786666.raw

198 <222> LOCATION: (11)
199 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
200 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
201 N-diisopropyl)phosphoramidite (6MAP, compound 11)
203 <400> SEQUENCE: 9

W--> 204 agaggtgtac naatgtggag a *see page 1* 21

207 <210> SEQ ID NO: 10
208 <211> LENGTH: 21
209 <212> TYPE: DNA
210 <213> ORGANISM: Artificial Sequence

212 <220> FEATURE:
213 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
214 HP24

216 <220> FEATURE:
217 <221> NAME/KEY: modified_base

218 <222> LOCATION: (11)
219 <223> OTHER INFORMATION: n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
220 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
221 N-diisopropyl)phosphoramidite (6MAP, compound 11)
223 <400> SEQUENCE: 10

W--> 224 agaggtgtac naatgtggag a 21

227 <210> SEQ ID NO: 11
228 <211> LENGTH: 21
229 <212> TYPE: DNA
230 <213> ORGANISM: Artificial Sequence

232 <220> FEATURE:
233 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
234 PTR31

236 <220> FEATURE:
237 <221> NAME/KEY: modified_base

238 <222> LOCATION: (7)
239 <223> OTHER INFORMATION: n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-
dimethoxytrityl)-

240 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
241 N-diisopropyl)phosphoramidite (DMAP, compound 12)
243 <400> SEQUENCE: 11

W--> 244 gtgtggnaaa tctctagcag t 21

247 <210> SEQ ID NO: 12
248 <211> LENGTH: 21
249 <212> TYPE: DNA
250 <213> ORGANISM: Artificial Sequence

252 <220> FEATURE:
253 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
254 PTR32

256 <220> FEATURE:
257 <221> NAME/KEY: modified_base

258 <222> LOCATION: (10)
259 <223> OTHER INFORMATION: n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-
dimethoxytrityl)-

260 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
261 N-diisopropyl)phosphoramidite (DMAP, compound 12)

RAW SEQUENCE LISTING
PATENT APPLICATION: US/09/786,666

DATE: 07/05/2001
TIME: 15:44:04

Input Set : A:\Nih-3511.app
Output Set: N:\CRF3\07032001\I786666.raw

263 <400> SEQUENCE: 12
W--> 264 gtgtggaaan tctctagcag t 21
267 <210> SEQ ID NO: 13
268 <211> LENGTH: 21
269 <212> TYPE: DNA
270 <213> ORGANISM: Artificial Sequence
272 <220> FEATURE:
273 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
274 PTR33
276 <220> FEATURE:
277 <221> NAME/KEY: modified_base
278 <222> LOCATION: (14)
279 <223> OTHER INFORMATION: n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-
dimethoxytrityl)-
280 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
281 N-diisopropyl)phosphoramidite (DMAP, compound 12)
283 <400> SEQUENCE: 13
W--> 284 gtgtggaaaa tctctnccag t 21
287 <210> SEQ ID NO: 14
288 <211> LENGTH: 21
289 <212> TYPE: DNA
290 <213> ORGANISM: Artificial Sequence
292 <220> FEATURE:
293 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
294 PTR34
296 <220> FEATURE:
297 <221> NAME/KEY: modified_base
298 <222> LOCATION: (7)
299 <223> OTHER INFORMATION: n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-
dimethoxytrityl)-
300 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
301 N-diisopropyl)phosphoramidite (DMAP, compound 12)
303 <400> SEQUENCE: 14
W--> 304 actgctnccag attttccaca c 21
307 <210> SEQ ID NO: 15
308 <211> LENGTH: 21
309 <212> TYPE: DNA
310 <213> ORGANISM: Artificial Sequence
312 <220> FEATURE:
313 <223> OTHER INFORMATION: Description of Artificial Sequence:oligonucleotide
314 PTR35
316 <220> FEATURE:
317 <221> NAME/KEY: modified_base
318 <222> LOCATION: (9)
319 <223> OTHER INFORMATION: n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-
dimethoxytrityl)-
320 beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
321 N-diisopropyl)phosphoramidite (DMAP, compound 12)
323 <400> SEQUENCE: 15
W--> 324 actgctnccag attttccaca c 21
327 <210> SEQ ID NO: 16
328 <211> LENGTH: 21

Please Note:

Use of n and/or Xaa have been detected in the Sequence Listing. Please review the
Sequence Listing to ensure that a corresponding explanation is presented in the <220> to
<223> fields of each sequence which presents at least one n or Xaa.

file://C:\Program Files\Nih-3511\I786666.raw

7/5/01

VERIFICATION SUMMARY

PATENT APPLICATION: US/09/786,666

DATE: 07/05/2001

TIME: 15:44:05

Input Set : A:\Nih-3511.app

Output Set: N:\CRF3\07032001\I786666.raw

L:15 M:271 C: Current Filing Date differs, Replaced Current Filing Date
L:44 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:1
L:64 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:2
L:84 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:3
L:104 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:4
L:124 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:5
L:144 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:6
L:164 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:7
L:184 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:8
L:204 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:9
L:224 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:10
L:244 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:11
L:264 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:12
L:284 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:13
L:304 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:14
L:324 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:15
L:344 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:16
L:364 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:17
L:384 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:18
L:404 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:19
L:424 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:20
L:444 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:21
L:464 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:22

PTO/PCT Rec'd 15 JUN 2001

PATENT
Attorney Docket No.: 015280-351100US

#3

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

HAWKINS *et al.*

Application No.: 09/786,666

Filed: March 07, 2001

For: PTERIDINE NUCLEOTIDE
ANALOGS

Examiner: Not yet assigned

Art Unit: Not yet assigned

COMMUNICATION UNDER

37 C.F.R. §§ 1.821-1.825

AND

PRELIMINARY AMENDMENT

Box SEQUENCE

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

In response to the Notification to Comply with Requirements for Patent Applications Containing Nucleotide Sequence and/or Amino Acid Sequence Disclosures, 37 C.F.R. §§ 1.821-1.825, that accompanied the Notification of Missing Requirements Under 35 U.S.C. 371 in the United States Designated/Elected Office (DO/EO/US) mailed 16 April, 2001, Applicants submit herewith the required paper copy and computer readable copy of the Sequence Listing. Please amend the specification in adherence with 37 C.F.R. §§ 1.821-1.825 as follows.

0978666-061501

In the Specification:

Please replace the paragraph (Table 2) beginning at page 39, line 1, with the following:

--Table 1

Relative quantum yields of oligonucleotides containing 6MAP or DMAP

SEQ ID NO:	Sequence (5' → 3')	w/6MAP	Q _{rel}
1	gtg tgg Faa atc tct agc agt	PTR21	0.010
2	gtg tgg aaa Ftc tct agc agt	PTR22	0.020
3	gtg tgg aaa atc tct Fgc agt	PTR23	0.018
4	act gct Fga gat ttt cca cac	PTR24	
5	act gct agF gat ttt cca cac	PTR25	0.011
6	act gct agc cFt ttt cca cac	PTR28	0.041
7	att cca caa Fgc cgt gtc a	HP21	0.010
8	aga ggt gtc cFc ctg tgg aga	HP22	<0.01
9	aga ggt gta cFa gtg tgg aga	HP23	0.012
10	aga ggt gta aFa atg tgg aga	HP24	<0.01
	Sequence (5' → 3')	w/DMAP	Q_{rel}
11	gtg tgg Faa atc tct agc agt	PTR31	0.023
12	gtg tgg aaa Ftc tct agc agt	PTR32	0.022
13	gtg tgg aaa atc tct Fgc agt	PTR33	0.01
14	act gct Fga gat ttt cca cac	PTR34	0.012
15	act gct agF gat ttt cca cac	PTR35	0.017
16	act gct aga gFt ttt cca cac	PTR36	0.019
17	act gct aga gat ttt ccF cac	PTR37	0.11
18	act gct agc cFt ttt cca cac	PTR38	0.11
19	att cca caa Fgc cgt gtc a	HP31	0.02
20	aga ggt gtc cFc ctg tgg aga	HP32	<0.01
21	aga ggt gta cFa gtg tgg aga	HP33	0.02
22	aga ggt gta aFa atg tgg aga	HP34	<0.01

--

Please insert the accompanying paper copy of the Sequence Listing, page numbers 1 to 8, at the end of the application.

09786666 OCT 2001

In the Claims:

Please amend claim 28 as follows:

28. (Once Amended) An oligonucleotide in accordance with claim 18, wherein said oligonucleotide is a member selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21 and SEQ ID NO:22.

REMARKS

A marked up copy of amended claim 28 is provided as an appendix entitled "**VERSION WITH MARKINGS TO SHOW CHANGES MADE.**" As a convenience to the Examiner, a complete set of the Claims, as amended herein, is also attached to this Amendment as an appendix entitled "**PENDING CLAIMS WITH ENTRY OF THE AMENDMENT**".


Applicants request entry of this amendment in adherence with 37 C.F.R. §§1.821 to 1.825. This amendment is accompanied by a floppy disk containing the above named sequences, SEQ ID NOS:1-22, in computer readable form, and a paper copy of the sequence information which has been printed from the floppy disk.

The information contained in the computer readable disk was prepared through the use of the software program "PatentIn" and is identical to that of the paper copy. This amendment contains no new matter.

09786666 "061501
T05T90" 9998260

If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 415-576-0200.

Respectfully submitted,


Kenneth A. Weber
Reg. No. 31,667

TOWNSEND and TOWNSEND and CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, California 94111-3834
Tel: (415) 576-0200
Fax: (415) 576-0300
JS:dmw

09/26/00 05:50:4

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Specification:

Paragraph (Table 2) beginning at line 1 of page 39 has been amended as follows:

Table 2

Relative quantum yields of oligonucleotides containing 6MAP or DMAP

SEQ ID NO:	Sequence (5' → 3')	w/6MAP	Q_{rel}
1	gtg tgg Faa atc tct agc agt	PTR21	0.010
2	gtg tgg aaa Ftc tct agc agt	PTR22	0.020
3	gtg tgg aaa atc tct Fgc agt	PTR23	0.018
4	act gct Fga gat ttt cca cac	PTR24	
5	act gct agF gat ttt cca cac	PTR25	0.011
6	act gct agc cFt ttt cca cac	PTR28	0.041
7	att cca caa Fgc cgt gtc a	HP21	0.010
8	aga ggt gtc cFc ctg tgg aga	HP22	<0.01
9	aga ggt gta cFa gtg tgg aga	HP23	0.012
10	aga ggt gta aFa atg tgg aga	HP24	<0.01
	Sequence (5' → 3')	w/DMAP	Q_{rel}
11	gtg tgg Faa atc tct agc agt	PTR31	0.023
12	gtg tgg aaa Ftc tct agc agt	PTR32	0.022
13	gtg tgg aaa atc tct Fgc agt	PTR33	0.01
14	act gct Fga gat ttt cca cac	PTR34	0.012
15	act gct agF gat ttt cca cac	PTR35	0.017
16	act gct aga gFt ttt cca cac	PTR36	0.019
17	act gct aga gat ttt ccF cac	PTR37	0.11
18	act gct agc cFt ttt cca cac	PTR38	0.11
19	att cca caa Fgc cgt gtc a	HP31	0.02
20	aga ggt gtc cFc ctg tgg aga	HP32	<0.01
21	aga ggt gta cFa gtg tgg aga	HP33	0.02
22	aga ggt gta aFa atg tgg aga	HP34	<0.01

09786666-064504
PTR21-064504

In the Claims:


Claim 28 has been amended as follows:

28. (Once Amended) An oligonucleotide in accordance with claim 18, wherein said oligonucleotide is a member selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21 and SEQ ID NO:22.

09786666-061E01

PENDING CLAIMS WITH ENTRY OF THE AMENDMENT

1. A compound of the formula:



wherein:


R¹ is a member selected from the group consisting of hydrogen and optionally substituted C₁-C₆-alkyl;

R² is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R³ is optionally substituted C₁-C₆ alkyl;

R⁴ is a member selected from the group consisting of hydrogen and L;

L is of the formula



wherein:

R⁵ is a member selected from the group consisting of hydrogen, hydroxyl, and substituted hydroxyl wherein the substituent is a protecting group;

R⁶ is a member selected from the group consisting of hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support, a hydroxyalkyl, and a hydroxyalkyl covalently bound to a solid support; and

R⁷ is a member selected from the group consisting of hydrogen, a phosphate, a triphosphate, and a protecting group;

with the proviso that R¹ and R⁴ are not simultaneously L.

2. A compound in accordance with claim 1, wherein R¹ is hydrogen; R² is a member selected from the group consisting of amino, mono-, and di-substituted amino wherein the substituents are members selected from the group consisting of benzoyl, isobutyryl, phthaloyl, di-n-butylaminomethylidene, dimethylaminomethylidene, p-nitrophenylethoxycarbonyl and dimethylaminomethylenamino;

R⁴ is L;

R⁵ is a member selected from the group consisting of hydrogen, hydroxyl, hydroxyl substituted with a member selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl, tetrahydropyran-1-yl, 4-methoxytetrahydropyran-4-yl, 1-(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl, t-butyl dimethylsilyl, p-nitrophenylethylsulfonyl, tetrahydropyranyl, 4-methoxytetrahydropyranyl, 2-nitrobenzyl, 9-phenylxanthen-9-yl and p-nitrophenylethyl;

R⁶ is a member selected from the group consisting of consisting of hydrogen, phosphoramidite, H-phosphonate, hemisuccinate, and hemisuccinate covalently bound to a solid support; and

R⁷ is a member selected from the group consisting of hydrogen, trityl, monomethoxytrityl, dimethoxytrityl, phthaloyl, di-n-butylaminomethylene, dimethylaminomethylidene and triphosphate.

3. A compound in accordance with claim 2, wherein R² is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

R⁵ is a member selected from the group consisting of hydrogen, hydroxyl and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyl dimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethylsulfonyl;

R⁶ is a member selected from the group consisting of hydrogen, β -cyanoethyl-N-diisopropyl phosphoramidite and a hemisuccinate covalently bound to controlled pore glass; and

R⁷ is a member selected from the group consisting of dimethoxytrityl, di-n-butylaminomethylene, and dimethylaminomethylidene.

09786666.051501

4. A compound in accordance with claim 2, wherein R² is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

R⁵ is a member selected from the group consisting of hydrogen and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethyl;

R⁶ is a member selected from the group consisting of hydrogen and β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is a member selected from the group consisting of hydrogen and dimethoxytrityl.

5. A compound in accordance with claim 2, wherein R² is a member selected from the group consisting of amino and dimethylaminomethylenamino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is dimethoxytrityl.

6. A compound in accordance with claim 2, wherein R² is amino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is dimethoxytrityl.

7. A compound in accordance with claim 2, wherein R² is amino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is hydrogen; and

R⁷ is hydrogen.

09786666-001501
T05T90" 99999760

8. A compound in accordance claim 2, wherein R² is dimethylaminomethylenamino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is dimethoxytrityl.

9. A compound in accordance with claim 2, wherein R² is amino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is a triphosphate.

10. A compound in accordance with claim 1, wherein;

R¹ is optionally substituted C₁-C₆ alkyl;

R² is a member selected from the group consisting of amino, mono-, and di-substituted amino wherein the substituent is a member selected from the group consisting of benzoyl, isobutyryl, phthaloyl, di-n-butylaminomethylidene, dimethylaminomethylidene, p-nitrophenylethoxycarbonyl and dimethylaminomethylenamino;

R³ is optionally substituted C₁-C₆ alkyl;

R⁴ is L;

R⁵ is a member selected from the group consisting of hydrogen, hydroxyl and hydroxyl substituted with a member selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl, tetrahydropyran-1-yl, 4-methoxytetrahydropyran-4-yl, 1-(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl, t-butyl dimethylsilyl, p-nitrophenylethylsulfonyl, tetrahydropyranyl, 4-methoxytetrahydropyranyl, 2-nitrobenzyl, 9-phenylxanthen-9-yl and p-nitrophenylethyl;

R⁶ is a member selected from the group consisting of hydrogen, H-phosphonate, phosphoramidite, hemisuccinate, and hemisuccinate covalently bound to a solid support; and

09786666-061501

R^7 is a member selected from the group consisting of hydrogen, trityl, monomethoxytrityl, dimethoxytrityl, phthaloyl, di-n-butylaminomethylene, and dimethylaminomethylidene.

11. A compound in accordance with claim 10 wherein R^1 is methyl;
 R^2 is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

R^3 is methyl;

R^5 is a member selected from the group consisting of hydrogen, hydroxyl and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyl dimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethylsulfonyl;

R^6 is a member selected from the group consisting of hydrogen, β -cyanoethyl-N-diisopropyl phosphoramidite and a hemisuccinate covalently bound to controlled pore glass; and

R^7 is a member selected from the group consisting of dimethoxytrityl, di-n-butylaminomethylene, and dimethylaminomethylidene.

12. A compound in accordance claim 10, wherein R^1 is methyl; R^2 is a member selected from the group consisting of amino and an amino group mono-substituted by a protecting group selected from the group consisting of di-n-butylaminomethylidene, p-nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

R^5 is a member selected from the group consisting of hydrogen and hydroxyl substituted with a member selected from the group consisting of dimethoxytrityl, tetrahydropyran-1-yl, t-butyl dimethylsilyl, 2-nitrobenzyl, and p-nitrophenylethylsulfonyl;

R^6 is a member selected from the group consisting of consisting of hydrogen and β -cyanoethyl-N-diisopropyl phosphoramidite; and

R^7 is a member selected from the group consisting of hydrogen and dimethoxytrityl.

0978666-061501

13. A compound in accordance with claim 10, wherein R¹ is methyl;
R² is a member selected from the group consisting of amino and
dimethylaminomethylenamino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is dimethoxytrityl.

14. A compound in accordance with claim 10, wherein R¹ is methyl;
R² is amino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is dimethoxytrityl.

15. A compound in accordance with claim 10, wherein R¹ is methyl;
R² is amino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is hydrogen; and

R⁷ is hydrogen.

16. A compound in accordance with claim 10, wherein R¹ is methyl; is
dimethylaminomethylenamino;

R³ is methyl;

R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

R⁷ is dimethoxytrityl.

17. A compound in accordance with claim 10, wherein R¹ is methyl;
R² is amino;

R³ is methyl;


R⁵ is hydrogen;

R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

09786666-051501

R^7 is a triphosphate.

18. An oligonucleotide comprising one or more nucleotide monomers, said monomers having the formula



wherein:


R^1 is a member selected from the group consisting of hydrogen and optionally substituted C_1 - C_6 -alkyl;

R^2 is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R^3 is optional substituted C_1 - C_6 alkyl;

R^4 is L;

L is of the formula



wherein:

R^5 is a member selected from the group consisting of hydrogen and hydroxyl;

R^6 is a member selected from the group consisting of hydrogen, a phosphate, a phosphate covalently attached to a nucleotide, a phosphate covalently attached to a nucleoside; a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide covalently bound to a solid support, and a hydroxyalkyl covalently bound to a solid support; and

R^7 is a member selected from the group consisting of hydrogen, a phosphate, a phosphate covalently attached to a nucleotide and a phosphate covalently attached to a nucleoside;

with the proviso that R¹ and R⁴ are not simultaneously L.

19. An oligonucleotide in accordance with claim 18, wherein:

R¹ is hydrogen;

R² is amino;

R³ is methyl;

R⁵ is hydrogen and hydroxyl;

R⁶ is hydrogen; and

R⁷ is a phosphate.

20. An oligonucleotide in accordance with claim 19, wherein:

R⁵ is hydrogen.

21. An oligonucleotide in accordance with claim 19 wherein:

R⁵ is hydroxyl.

22. An oligonucleotide in accordance with claim 18, wherein:

R¹ is optionally substituted C₁-C₆-alkyl;

R² is amino;

R³ is methyl;

R⁵ is hydrogen and hydroxyl;

R⁶ is hydrogen; and

R⁷ is a phosphate.

23. An oligonucleotide in accordance with claim 22, wherein

R¹ is methyl and

R⁵ is hydrogen.

24. An oligonucleotide in accordance with claim 22, wherein

R¹ is methyl and

R⁵ is hydroxyl.

25. An oligonucleotide in accordance with claim 18, wherein said nucleotide monomers are at the 3' end of said oligonucleotide.

0978666-061501
T 05T90 99999260


26. An oligonucleotide in accordance with claim 18, wherein said nucleotide monomers are at the 5' end of said oligonucleotide.

27. An oligonucleotide in accordance with claim 18, wherein said nucleotide monomers are surrounded by 1 to 10 pyrimidine monomers.

28. An oligonucleotide in accordance with claim 18, wherein said oligonucleotide is a member selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21 and SEQ ID NO:22.

29. A method of detecting the presence, absence, or quantity of a target nucleic acid, said method comprising the steps of:

a) contacting said target nucleic acid with a nucleic acid probe wherein said nucleic acid probe comprises compound of the formula:



wherein:


R¹ is a member selected from the group consisting of hydrogen and optionally substituted C₁-C₆-alkyl;

R² is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R³ is optionally substituted C₁-C₆ alkyl;

R⁴ is L;

L is of the formula



wherein:

R⁵ is a member selected from the group consisting of hydrogen and hydroxyl;

R⁶ is a member selected from the group consisting of hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support; and

R⁷ is phosphate;

with the proviso that R¹ and R⁴ are not simultaneously L; located in said probe such that, when said probe hybridizes to said target nucleic acid said compound is in a loop that does not participate in complementary base pairing with a nucleotide of said target nucleic acid; and

b) detecting the fluorescence produced by said fluorescent nucleotide when said probe forms a hybrid duplex with said target nucleic acid.

30. A method of claim 29, wherein said loop ranges in length from about 1 to about 100 nucleotides when said probe hybridizes to said target nucleic acid.

31. A method of claim 29, wherein said loop is an insertion in said nucleic acid probe which is otherwise complementary to said target nucleic acid or to a contiguous subsequence of said target nucleic acid.

32. A method of claim 31, wherein said insertion is three nucleotides in length and comprises two nucleotides each adjacent to said compound.

33. A method of claim 32, wherein at least one nucleotide adjacent to said compound is a purine.

34. A method of claim 33, wherein at least one nucleotide adjacent to said compound is an adenosine.

35. A method of claim 32, wherein at least one nucleotide adjacent to said compound is a pyrimidine.

09786666-001501

36. A method of claim 35, wherein at least one nucleotide adjacent to said compound is a cytosine.

37. A method of claim 34, wherein said compound is bordered by at least two adjacent purines in both the 5' and 3' direction.

38. A method of claim 37, wherein said adjacent purines are adenosine.

39. A method of claim 31, wherein said insertion is said compound.

40. A method of claim 31, wherein said insertion is self-complementary and forms a hairpin wherein said compound is present in the loop of said hairpin and does not participate in complementary base pairing.

41. A method of claim 29, wherein the nucleotides comprising said loop are selected such that they are not complementary to the corresponding nucleotides of the target nucleic acid when said probe is hybridized to said target nucleic acid and wherein said probe is complementary to at least two non-contiguous subsequences of said target nucleic acid.

42. A method of claim 29, wherein said fluorescent nucleotide is present in a terminal subsequence of said nucleic acid probe wherein said terminal subsequence does not hybridize to said target nucleic acid when the remainder of said nucleic acid probe hybridizes to said target nucleic acid.

43. A method of claim 42, wherein said terminal subsequence forms a terminal hairpin by hybridization with a second subsequence of said probe such that said fluorescent nucleotide is present in a loop of said hairpin and does not participate in complementary base pairing.

44. A method of claim 29, wherein said detecting comprises detecting an increase in fluorescence of said fluorescent nucleotide when said probe forms a hybrid duplex with said target nucleic acid.

0978666-051504
PATENT

45. A kit for the detection of nucleic acid-nucleic acid interactions comprising a container, said container containing a compound in accordance with claim 1, and instructions for use.

SF 1233603 v1

09/28/2011 10:51:50

SEQUENCE LISTING

<110> Hawkins, Mary E.
Pfleiderer, Wolfgang
Balis, Frank
The Government of the United States of America
as represented by The Secretary of the
Department of Health and Human Services

<120> Pteridine Nucleotide Analogs

<130> 015280-351100US

<140> US 09/786,666
<141> 2001-03-07

<150> US 60/099,487
<151> 1998-09-08

<150> WO PCT/US99/20541
<151> 1999-09-07

<160> 22

<170> PatentIn Ver. 2.1

<210> 1
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR21

<220>
<221> modified_base
<222> (7)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 1
gtgtggnaaa tctctagcag t 21

<210> 2
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR22

<220>
<221> modified_base
<222> (10)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 2
gtgtggaaan tctctagcag t

21

<210> 3
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR23

<220>
<221> modified_base
<222> (16)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 3
gtgtggaaaa tctctngcag t

21

<210> 4
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR24

<220>
<221> modified_base
<222> (7)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 4
actgctngag attttccaca c

21

<210> 5
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR25

<220>
<221> modified_base
<222> (9)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 5
actgctagn g attttccaca c 21

<210> 6
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR28

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 6
actgctagcc nttttccaca c 21

<210> 7
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP21

<220>
<221> modified_base
<222> (10)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 7
attccacaan gccgtgtca 19

<210> 8
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP22

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 8
agaggtgtcc ncctgtggag a

21

<210> 9
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP23

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 9
agaggtgtac nagtgtggag a

21

<210> 10
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP24

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (6MAP, compound 11)

<400> 10
agaggtgtaa naatgtggag a

21

<210> 11
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR31

<220>
<221> modified_base
<222> (7)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 11
gtgtggnaaa tctctagcag t

21

<210> 12
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR32

<220>
<221> modified_base
<222> (10)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 12
gtgtggaaan tctctagcag t

21

<210> 13
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR33

<220>
<221> modified_base
<222> (14)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 13
gtgtggaaaa tctctngcag t

21

<210> 14
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR34

<220>
<221> modified_base
<222> (7)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 14
actgctngag attttccaca c 21

<210> 15
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR35

<220>
<221> modified_base
<222> (9)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 15
actgctagng attttccaca c 21

<210> 16
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR36

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 16
actgctagag nttttccaca c 21

<210> 17
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR37

<220>
<221> modified_base
<222> (18)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 17
actgctagag attttccnca c

21

<210> 18
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
PTR38

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 18
actgctagcc nttttccaca c

21

<210> 19
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP31

<220>
<221> modified_base
<222> (10)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 19
attccacaan gccgtgtca

19

<210> 20
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP32

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 20
agaggtgtcc ncctgtggag a

21

<210> 21
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP33

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 21
agaggtgtac nagtgtggag a

21

<210> 22
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide
HP34

<220>
<221> modified_base
<222> (11)
<223> n = 4-amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-
beta-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-beta-cyanoethyl
N-diisopropyl)phosphoramidite (DMAP, compound 12)

<400> 22
agaggtgtaa naatgtggag a

21

PTERIDINE NUCLEOTIDE ANALOGS

RELATED APPLICATIONS

The application claims priority to U. S. Provisional Patent Application No. 60/099,487 filed September 8, 1998, the teachings of which are incorporated herein by reference in their entirety for all purposes.

BACKGROUND OF THE INVENTION

Fluorescence is a useful analytical technique for studying nucleic acid/nucleic acid and protein/nucleic acid interactions. In particular, fluorescent nucleosides and their analogs can be used to probe the physical and chemical environment of macromolecules to which they are covalently bonded or electrostatically attached. Various physical and chemical environmental changes can be assessed using fluorescence such as, changes in pH, interaction with other molecules, or alteration in tertiary structure. These changes can manifest differences in fluorescence intensity, the lifetime of fluorescence emission or fluorescence depolarization. In addition, there can be a shift in the emission, excitation, or absorption spectra of the fluorescent nucleoside depending on its environment. By measuring changes in fluorescence, useful information relating to macromolecule interactions, nucleic acid hybridization and enzymatic reactions can be obtained.

Fluorescent oligonucleotides find additional uses in molecular biology as probes for screening genomic and complementary DNA libraries, as primers for DNA synthesis, sequencing, and amplification reactions. Oligonucleotide probes have also proven useful for assaying *in vitro* gene expression using techniques of *in situ* hybridization. Recent improvements in DNA sequencing methods, fluorescent labels, and detection systems have dramatically increased the use of fluorescently labeled oligonucleotides in all of the foregoing applications. Typically, oligonucleotides are labeled with a fluorescent marker, either directly through a covalent linkage (e.g., a carbon linker), intercalation, or indirectly whereby the oligonucleotide is bound to a molecule such as biotin or dioxigenin, which is subsequently coupled to a fluorescently labeled binding moiety (e.g., streptavidin or a labeled monoclonal antibody).


These fluorescent labeling systems, however, suffer the disadvantage that the fluorescent complexes and their binding moieties are relatively large. The presence of

large fluorescent labels and associated linkers can alter the mobility of the oligonucleotide, either through a gel as in sequencing, or through various compartments of a cell. Also, the means of attachment, typically through a 6-carbon linker, positions the probe at some distance from the other bases and allows movement of the probe in ways unrelated to the movement of the oligonucleotide. This can distort fluorescent depolarization measurements.

Ideally, a fluorescent nucleoside analog should closely resemble the naturally occurring purine or pyrimidine base structure. In particular, the probe should be attached to the oligonucleotide through the native deoxyribose chain. This keeps the probe aligned with other bases in the oligonucleotide and allows it to move in a more native-like manner. The analog should especially possess similar hydrogen bonding interactions. One type of nucleoside analog is a furanosyl pteridine derivative. Pteridines are a class of bicyclic planar compounds, some of which are highly fluorescent and are structurally similar to purines (*see*, Figure 1). In fact, the fluorescence of many pteridine nucleoside analogs is known. U.S. Patent No. 5,525,711, herein incorporated by reference, discloses pteridine nucleotide analogs as fluorescent DNA probes.

SUMMARY OF THE INVENTION

The present invention provides fluorescent nucleoside analogs that closely resemble the naturally occurring purine base structure. These pteridine nucleotides are much more stable and possess higher quantum yields than prior art compounds. As such, the present invention provides pteridine nucleotides of Formula I




I

In Formula I, R¹ is a functional group including, but not limited to, hydrogen and optionally substituted C₁-C₆-alkyl.

In Formula I, R² is a functional group including, but not limited to, amino and mono- or di-substituted amino wherein the substituent(s) is a protecting group.

In Formula I, R³ is a functional group including, but not limited to, optionally substituted C₁-C₆-alkyl.

In Formula I, R^4 is a functional group including, but not limited to, hydrogen and a compound having formula L.



L

In Formula L, R^5 is a functional group including, but not limited to, hydrogen, hydroxyl and substituted hydroxyl wherein the substituent is a protecting group.

In Formula L, R^6 is a functional group including, but not limited to, hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support.

In Formula L, R^7 is a functional group including, but not limited to, hydrogen, a phosphate, a triphosphate, and a protecting group. In addition, R^1 and R^4 are not simultaneously L.

Compounds of Formula I are highly fluorescent under normal physiological conditions, and suitable for use in the chemical synthesis of oligonucleotides.

In another embodiment, the present invention relates to oligonucleotides that incorporate these pteridine nucleotides.

In yet another embodiment, the present invention relates to pteridine nucleotide triphosphates that may be utilized in various nucleic acid amplification processes. When used in a nucleic acid amplification process, the pteridine nucleotide triphosphates are directly incorporated into the amplified sequence rendering it fluorescent.

In still yet another embodiment, this invention relates to methods of detecting the presence, absence, or quantity of a target nucleic acid. The methods involve probing the target nucleic acid with a nucleic acid probe identical in sequence to the target sequence with the addition of the pteridine probe that does not have a pairing partner. When annealing occurs, the pteridine probe is squeezed out of the base stacking

into a loop. This removes the pteridine probe from the quenching effects of base stacking and yields an increase in fluorescence intensity.

In one preferred embodiment, the loop in the above probe ranges in length from about 1 to about 100 nucleotides when the probe hybridizes to the target nucleic acid. In particularly preferred probes, the loop is an insertion in the nucleic acid probe that is otherwise complementary to the target nucleic acid or to a contiguous subsequence of the target nucleic acid. In some preferred embodiments, the insertion is three nucleotides in length and which two nucleotides are each adjacent to the fluorescent nucleotide. In particularly preferred embodiments, at least one nucleotide adjacent to the fluorescent nucleotide is a purine (*e.g.*, adenosine), and in still more preferred embodiments, the fluorescent nucleotide is bordered by at least two adjacent purines (*e.g.*, adenosine) in both the 5' and 3' direction. In a most preferred embodiment, the insertion is a single base insertion; a pteridine nucleotide of Formula I.

In yet another embodiment, the insertion is self-complementary and forms a hairpin in which the fluorescent pteridine nucleotide is present in the loop of the hairpin and does not participate in complementary base pairing. The nucleotides comprising the loop can be selected such that they are not complementary to the corresponding nucleotides of the target nucleic acid when the probe is hybridized to the target nucleic acid and where the probe is complementary to at least two non-contiguous subsequences of the target nucleic acid.

In still yet another embodiment, the invention also provides kits for performing nucleic acid amplifications or for detecting the presence absence or quantity of a nucleic acid in a sample. The kits comprise a container containing any of the probes or label oligonucleotide having a compound of Formula I described herein. The kit can further comprise, a buffer, and/or any of the other reagents useful for practicing the method to which the kit is directed. These and other embodiments of the present invention will be described in detail below.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the numbering system of a pteridine molecule.

Figure 2 illustrates a synthetic scheme of compounds of Formula I.

Figure 3 illustrates excitation and emission scans of 6MAP (solid line – higher peaks) and DMAP (dashed line – lower peaks). Samples were measured in 10 mM Tris pH 7.5 at room temperature.

Figure 4 illustrates effects of P1 nuclease digestion on oligonucleotides containing 6MAP. Reaction mixtures were incubated overnight at 37°C.

Figure 5 illustrates effects of P1 nuclease digestion on oligonucleotides containing DMAP. Reaction mixtures were incubated overnight at 37°C

Figure 6 illustrates P1 nuclease digestion as it occurs in the time-based acquisition mode. The blank rate contains all the same components with the exception of P1 nuclease.

DETAILED DESCRIPTION OF THE INVENTION

I. Glossary

The term "C₁-C₆-alkyl" denotes branched or unbranched hydrocarbon chains, such as, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, octa-decyl and 2-methylpentyl. These groups can be optionally substituted with one or more functional groups which are attached commonly to such chains, such as, hydroxyl, bromo, fluoro, chloro, iodo, mercapto or thio, cyano, alkylthio, heterocyclyl, aryl, heteroaryl, carboxyl, carbalkoyl, alkyl, alkenyl, nitro, amino, alkoxyl, amido, and the like to form alkyl groups such as trifluoromethyl, 3-hydroxyhexyl, 2-carboxypropyl, 2-fluoroethyl, carboxymethyl, cyanobutyl and the like.

The term "oligonucleotide" refers to a molecule comprised of two or more deoxyribonucleotides, ribonucleotides, modified ribonucleotides, modified deoxyribonucleotides, ribonucleotide analogs, deoxyribonucleotide analogs, peptide nucleic acids, pteridine derivatives of the present invention, and other chemically modified nucleic acids. The exact size of an oligonucleotide depends on many factors and the ultimate function or use of the oligonucleotide. Generally, chemically synthesized oligonucleotides range in length from 2 to 500 bases, although, it is well known that oligonucleotides may be ligated together to provide longer sequences. As used herein, the term "oligonucleotide" also encompasses these longer sequences. It is also recognized that double-stranded polynucleotides may be created by hybridization with a complementary sequence or enzymatically through primer extension. The term oligonucleotide as used in this application encompasses both single and double-stranded oligonucleotide.

The term "solid support" refers to a solid material that is functionalized to permit the coupling of a monomer used in polynucleotide synthesis. The solid support is

typically coupled to a nucleoside monomer through a covalent linkage to the 3'-carbon on the furanose. Solid support materials typically are unreactive during the polynucleotide synthesis and simply provide a substratum to anchor the growing polynucleotide. Solid support materials include, but are not limited to, polacryloylmorpholide, silica, controlled pore glass (CPG), polystyrene, polystyrene/latex, and carboxyl modified Teflon.

The term "cleavage" in reference to solid phase oligonucleotide synthesis refers to the breaking of the bond that binds an oligonucleotide to a solid support. Typically, cleavage involves hydrolysis of a succinate ester bond between the 3'-hydroxyl of an attached oligonucleotide and the solid support.

The term "pteridine nucleotide" or "pteridine monomer" is used herein to refer to the furanosyl pteridine derivatives of the present invention with a 3'-phosphate group. It is recognized that properly speaking the furanosyl pteridine derivatives are not nucleotides as the pteridine is neither a purine nor a pyrimidine. However, because the furanosyl pteridine derivatives are structurally analogous to purine nucleotides, and the furanosyl pteridines of this invention are used in the same manner as nucleotides both will be referred to as nucleotides. As used herein, the pteridine nucleotide or pteridine monomer may be fully protected for use in polynucleotide synthesis or it may be deprotected when used as a triphosphate or when incorporated into an oligonucleotide

The term "nucleotide monomer" as used herein refers to pteridine nucleotides, the "standard" nucleotides; adenosine, guanosine, cytidine, thymidine, and uracil, or derivatives of these nucleotides. Such derivatives include, but are not limited to, inosine, 5-bromodeoxycytidine, 5-bromo-deoxyuridine, N⁶-methyl-deoxyadenosine and 5-methyl-deoxycytidine.

As used herein, the term "protecting group" refers to a group that is joined to or substituted for a reactive group (e.g., a hydroxyl or an amine) on a molecule. The protecting group is chosen to prevent reaction of the particular radical during one or more steps of a chemical reaction. Generally the particular protecting group is chosen so as to permit removal at a later time to restore the reactive group without altering other reactive groups present in the molecule. The choice of a protecting group is a function of the particular radical to be protected and the compounds to which it will be exposed. The selection of protecting groups is well known to those of skill in the art. See, for example Greene *et al.*, *Protective Groups in Organic Synthesis*, 2nd ed., John Wiley & Sons, Inc. Somerset, N.J. (1991).

As used herein, the term "protected amine" refers to an amine that has been reacted with an amino protecting group. An amino protecting group prevents reaction of the amide function during either the synthesis of the derivatized pteridine nucleoside or during the chemical synthesis of DNA or RNA using that nucleotide. The amino protecting group can be removed at a later time to restore the amino group without altering other reactive groups present in the molecule. For example, the exocyclic amine may be reacted with dimethylformamide diethylacetal to form the dimethylaminomethylenamino function. Amino protecting groups generally include carbamates, benzyl radicals, imidates, and others known to those of skill in the art. Preferred amino protecting groups include, but are not limited to, p-nitrophenylethoxycarbonyl or dimethylaminomethylenamino.

The term "coupling" is generally used in DNA synthesis to refer to the joining of one nucleotide monomer to another nucleotide monomer or to the 5' terminal of an oligonucleotide. The coupling is generally accomplished by the formation of a phosphodiester linkage from the 3'-phosphate of one nucleotide monomer to the 5'-hydroxyl of a second monomer or oligonucleotide. Coupling is also used to refer to the joining of an initial nucleoside to a solid support.

The term "label oligonucleotide", as used herein, refers to an oligonucleotide incorporating one or more fluorescent nucleotide analogues. The fluorescence activity of the nucleotide analogue(s) can be quenched partially or to a non-detectable level when the label oligonucleotide achieves a substantially linear conformation (*i.e.*, the constituent bases, more particularly the fluorescent nucleotide(s), participate in normal base stacking). Preferred label oligonucleotide of this invention are capable of achieving a conformation, when hybridized to themselves or another nucleic acid or when bound by a nucleic acid binding protein, in which the quench (reduction of fluorescence intensity of the fluorescent nucleotide(s)) is diminished or eliminated resulting in a label oligonucleotide having increased fluorescence when present in that conformation. The label oligonucleotides of this invention are distinguished from labeled oligonucleotides wherein a label is attached. The labeled oligonucleotide can of course be attached to (labeled with) a label oligonucleotide of the present invention either directly through a phosphodiester linkage or indirectly through a linker.

The terms "target nucleic acid" or "target oligonucleotide" refer to the nucleic acid sequence or nucleic acid subsequence that is to be detected using one or

more label oligonucleotide of this invention. The label oligonucleotides typically hybridize to all or a part of the target nucleic acid under stringent conditions.

The term "corresponding nucleotide", is used to refer to the position of a nucleotide in a first nucleic acid by reference to a second nucleic acid. Thus, a
5 corresponding nucleotide refers to a nucleotide that it is positionally located opposite to a base where neighboring bases are all hybridized pairs

The term "probe" refers to the nucleic acid sequence or nucleic acid subsequence that is used to detect a target nucleic acid. In an especially preferred embodiment, a probe is a label oligonucleotide of this invention.

10 "Subsequence" refers to a sequence of nucleic acids that comprise a part of a longer sequence of nucleic acids.

Hybridization refers to the specific binding of two nucleic acids through complementary base pairing. Hybridization typically involves the formation of hydrogen bonds between nucleotides in one nucleic acid and their corresponding nucleotides in the
15 second nucleic acid.

"Bind(s) substantially" refers to complementary hybridization between a probe nucleic acid and a target nucleic acid and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired detection of the target polynucleotide sequence.

20 The phrase "hybridizing specifically to", refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence or subsequence under stringent conditions when that sequence is present in a complex mixture (*e.g.*, total cellular DNA or RNA).

The term "stringent conditions" refers to conditions under which a probe
25 will hybridize to its target subsequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined
30 ionic strength, pH, and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. (As the target sequences are generally present in excess, at T_m , 50% of the probes are occupied at equilibrium) Typically, stringent conditions will be those in which the salt concentration is at most about 0.01 to 1.0 M Na ion concentration (or other salts) at pH

7.0 to 8.3 and the temperature is at least about 30°C for short probes (*e.g.*, 10 to 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

The term complementary base pair refers to a pair of bases (nucleotides) each in a separate nucleic acid in which each base of the pair is hydrogen bonded to the other. A "classical" (Watson-Crick) base pair always contains one purine and one pyrimidine; adenine pairs specifically with thymine (A-T), guanine with cytosine (G-C), uracil with adenine (U-A). The two bases in a classical base pair are said to be complementary to each other.

A "nucleic acid amplification mixture" refers to the reaction mixture used to amplify a nucleic acid. The amplification may be by any method including but not limited to PCR, long range PCR, ligase chain reaction, self-sustained sequence replication, and the like. Typical nucleic acid amplification mixtures (*e.g.*, PCR reaction mixture) include a nucleic acid template that is to be amplified, a nucleic acid polymerase, nucleic acid primer sequence(s), and nucleotide triphosphates, and a buffer containing all of the ion species required for the amplification reaction.

II. *Fluorescent Nucleosides*

In certain aspects, the compounds of this invention are fluorescent nucleosides that can be used in a great variety of biological and physical chemistry applications. They can be used for instance, as fluorescent labels to label almost any biological molecule, as well as to probe the physical and chemical environment of macromolecules. The nucleosides of this invention can also be derivitized as nucleotide triphosphates that can then be utilized as monomers for DNA synthesis. In one preferred embodiment, the pteridine nucleotides of Formula I are derivitized for DNA synthesis by protecting the reactive exocyclic amine, for example, R² can be a dimethylaminomethylenamino group, the 3'-hydroxyl can be derivitized as a phosphoramidite group and R⁷ can be derivitized as a dimethoxytrityl group.

Moreover, the compounds of Formula I can be used as fluorescent labels. In certain embodiments, the pteridine nucleotides can be linked through the 5'-hydroxyl, the 3'-phosphate, or the 2'-hydroxyl (in the case of a ribofuranose) directly, or through a linker, to the composition it is desired to label. Such labeled compositions can include, but are not limited to, biological molecules such as antibodies, ligands, lipids, polysaccharides, cell surface receptors, and enzymes.


In addition, the pteridine nucleotide triphosphates of Formula I can be used in DNA amplification techniques such as, the polymerase chain reaction. For instance, a pair of PCR primers can be chosen that are complementary to the DNA sequences flanking a DNA sequence of interest. The PCR reaction mixture will contain one or more species of nucleotide triphosphates of Formula I. If the proper target sequences are present in the sample, the DNA sequence between the primers will be amplified. This amplified DNA sequence will then contain the fluorescent pteridine nucleotide triphosphates of the present invention.

Moreover, the compounds of this invention can be used as hybridization probes to detect the presence, absence, or quantity of a target nucleic acid. In one embodiment, this involves contacting the target nucleic acid with a nucleic acid probe where the nucleic acid probe comprises a compound of Formula I located in the probe. When the probe hybridizes to the target nucleic acid, the compound of Formula I is forced out of base stacking which relieves fluorescence quenching caused by base stacking. This results in multi-fold increases in fluorescence intensity. These as well as other applications and uses will be described in detail below.

III. Preferred Fluorescent Nucleosides and Their Synthesis

There is a great need for fluorescent nucleosides that are purine analogs, especially adenosine analogs, which are more stable and possess higher fluorescent quantum yields than the prior art compounds. The inventors have found that the compounds of Formula (I) are more stable and possess greater quantum yields than the compounds of the prior art.

As such, the present invention relates to compounds of Formula (I)



wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and L have been defined above.

In Formula I, certain compounds are preferred. In one embodiment, compounds where R^1 is hydrogen; R^2 is amino or dimethylaminomethylenamino; R^3 is methyl; R^4 is L; R^5 is hydrogen; R^6 is hydrogen or β -cyanoethyl-N-diisopropyl phosphoramidite; and R^7 is hydrogen, dimethoxytrityl or a triphosphate are preferred.

Compounds of Formula I are more stable and have a higher fluorescence quantum yield than structurally similar compounds.

In another embodiment, compounds where R¹ is C₁-C₆ alkyl, especially methyl; R² is amino or dimethylaminomethylenamino; R³ is methyl; R⁴ is L; R⁵ is hydrogen; R⁶ is hydrogen or β -cyanoethyl-N-diisopropyl phosphoramidite; and R⁷ is hydrogen, dimethoxytrityl or a triphosphate are preferred.

The synthesis of compounds of Formula I is accomplished by reacting a protected pteridine derivative with a chlorofuranose having its 3'- and 5'-hydroxyls protected as their 4-chlorobenzoyl or paratoluoyl esters to produce a pteridine nucleoside. Following coupling, the protecting groups can be removed and the 5'-hydroxyl converted to its dimethoxytrityl ether. Subsequent conversion of the 3'-hydroxyl to the H-phosphonate, phosphoramidite, or hemisuccinate provides the compounds of Formula I

More specifically, with reference to Figure 2, 4,5,6-triaminopyrimidine (1), (*see*, J. Baddiley *et al.*, *J. Chem. Soc.* 386 (1943)) and ethyl pyruvate are heated in glacial acetic acid for 2 hours. After cooling the precipitate is collected and washed with water and purified by recrystallization using a suitable solvent system, such as DMF/H₂O, to yield 4-amino-6-methyl-7(8H)-pteridone (3) (*see*, D. Soll *et al.*, *Chem. Ber.* 96:2977 (1963)). Compounds wherein R¹ is an methyl group can be generated similarly by starting with 4,5,6-triamino-2-methylpyrimidine (2).

To covalently attach the furanose ring, compound (3) or (4) and DBU (1,8-diaza-bicyclo[5.4.0]-undec-7-ene) are stirred in an anhydrous solvent, such as acetonitrile, and then 2-deoxy-3,5-di-O-p-chlorobenzoyl- α -D-ribofuranosyl chloride (J. J. Fox *et al.*, *J. Am. Chem. Soc.* 83:4066 (1961)) is added. Stirring is continued for 2 hours which generates compounds (5) and (6), respectively. Subsequent removal of the chlorobenzoyl protecting group using anhydrous methanol will yield compounds (7) or (8), respectively.

The 5'-hydroxyl of the nucleoside can then be blocked with a protecting group (preferably dimethoxytrityl). Means of coupling protecting groups are well known to those of skill in the art. This is accomplished by reaction of the nucleoside with dimethoxytrityl chloride in dry pyridine. Other protocols are generally known to those of skill in the art. More specifically, with reference to Figure 2, compound (7) or (8) is reacted with 4,4'-dimethoxytrityl chloride in a suitable solvent, such as pyridine, to yield compound (9) or (10), respectively.

Where a protected amine is desired in the product, it can be introduced at any of several stages. For example, the starting pteridine can contain an amine substituent that is protected prior to further manipulation. Alternatively, an amine can be introduced at a later stage by conversion of an oxo moiety to a thione followed by amination with ammonia. Yet another method for introducing an amine uses a starting pteridine bearing a methylthio substituent in the 2-position. After coupling with the desired chlorofuranose, the protecting groups are removed and the methylthio group is displaced with ammonia.

The 3'-hydroxyl of the pteridine nucleoside can be converted to its respective hemisuccinate, phosphoramidite, H-phosphonate, or triphosphate using methods known to those of skill in the art. For example, conversion of the nucleoside 3'-hydroxyl to a hemisuccinate can be accomplished by reaction with succinic anhydride. Means of converting a nucleoside to a phosphoramidite are also well known to those of skill in the art.

More specifically, with reference to Figure 2, 4-amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-ribofuranosyl]-7(8H)-pteridone-3'-O-(2- β -cyanoethyl N-diisopropyl)phosphoramidite compound (11) or its methyl analog compound (12), can be synthesized starting from compound (9) or (10) respectively and bis-(N,N-diisopropylamino)-2- β -cyanoethoxyphosphane and 1H-tetrazole.

Similarly, means of converting a nucleoside to an H-phosphonate are also well known to those of skill in the art. In one approach, phosphorous (III) trichloride derivatives are used to directly phosphitylate the 3'-hydroxyl of the nucleoside. More specifically, phosphorous (III) triimidazolidine can be used to phosphitylate the 3'-hydroxyl. This method is described in detail by Garegg *et al.*, *Chemica Scripta* 25:280-282 (1985) and by Tocik *et al.*, *Nucleic Acids Res.* 18:193 (1987). Similarly, the use of tris-(1,1,1,3,3,3-hexafluoro-2-propyl) phosphite to produce ribonucleoside-H-phosphonates is described by Sakatsume *et al.*, *Nucleic Acids Res.* 17:3689-3697 (1989). The use of the same reagent to produce deoxynucleoside-H-phosphonates is described by Sakatsume *et al.*, *Nucleic Acids Res.* 18:3327-3331 (1990). Other approaches to the derivatization of the 3'-hydroxyl to produce H-phosphonates may be found in Sekine *et al.*, *J. Org. Chem.* 47:571-573 (1982); Marugg *et al.*, *Tetrahedron Lett.* 23:2661-2664 (1986), and Pon *et al.*, *Tetrahedron Lett.* 26:2525-2528 (1985).

Derivatization of the 5'-hydroxyl as a triphosphate can be accomplished by a number of means known to those of skill in the art. Where the pteridine nucleoside has

sufficient structural similarity to native nucleotides to act as an enzymatic substrate, the monophosphate may be synthesized chemically as described below and then enzymatically converted to the diphosphate and then to the triphosphate using the appropriate nucleotide monophosphate and diphosphate kinases respectively.


5 Alternatively, the nucleoside can be chemically derivatized as the triphosphate. This may be accomplished by reacting the nucleoside with trimethyl phosphate and POCl_3 and then adding a triethylammonium bicarbonate buffer to form the nucleotide monophosphate which can then be purified chromatographically. The nucleotide monophosphate is then activated using carbonyldiimidazole and coupled with
10 tributylammonium pyrophosphate to form the nucleotide triphosphate. The nucleotide triphosphate can then be precipitated as a sodium salt.

IV. *Oligonucleotide Synthesis*

As previously discussed, the pteridine derivatives of the present invention
15 are structurally analogous to adenosine. In certain aspects of the present invention, the compounds of Formula I are incorporated into an oligonucleotide, where they act as fluorescent tags. Because the compounds of Formula I are structurally analogous to the naturally occurring base, they do not alter the physical and chemical properties of the oligonucleotide as severely as currently available fluorescent tags. In some cases, the
20 perturbations are so minimal as to allow the oligonucleotide to act as an enzyme substrate permitting the enzyme catalyzed reaction to occur even when the substitution has been made at a site known to be critical for the enzyme function. Thus the oligonucleotides of this invention are particularly useful in the investigation of DNA-protein interactions.

In certain aspects, the compounds of Formula I can be at the terminal
25 position of the oligonucleotide or, alternatively, an internal position. The oligonucleotide can contain one or more nucleotide monomers of Formula I. The nucleotide monomers can be the same or different. The compounds can be adjacent to one another or spatially separated. If the oligonucleotide contains more than one compound of Formula I their arrangement can be random or quite specific, *e.g.*, being next only to purines or to
30 pyrimidines etc.

As such, this invention relates to an oligonucleotide comprising one or more nucleotide monomers Formula I



wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and L have been defined above.

Compounds of Formula I wherein R^6 is β -cyanoethyl, N-diisopropyl phosphoramidite are preferred as oligonucleotide synthesis monomers. These compounds can generally be utilized in most commercial DNA synthesizers without modification of the synthesis protocol. However, where large scale synthesis is desired, or where it is desirable to incorporate sulfur groups or other modifications in the phosphate linkages, compounds of Formula I wherein R^6 is H-phosphonate are preferred as synthesis reagents. The synthesis and use of other phosphorous containing derivatives suitable for oligonucleotide synthesis is well known to those of skill in the art. These include, but are not limited to, a methyl phosphonate, a phosphorothioate, and a phosphotriester. Thus, in one preferred embodiment, the pteridine nucleotides of Formula I are derivatized and protected for use as reagents in the synthesis of oligonucleotide. In particular, the reactive exocyclic amines are protected, for example R^2 is dimethylaminomethylenamino, the 3'-hydroxyl is derivatized as an H-phosphonate, or as a phosphoramidite and R^7 is dimethoxytrityl. Table 2 sets forth preferred oligonucleotides of the present invention.

The oligonucleotide of the present invention can be synthesized in solid phase or in solution. Generally, solid phase synthesis is preferred. Detailed descriptions of the procedures for solid phase synthesis of oligonucleotide by phosphite-triester, phosphotriester, and H-phosphonate chemistries are widely available. See, for example, Itakura, U.S. Pat. No. 4,401,796; Caruthers *et al.*, U.S. Pat. Nos. 4,458,066 and 4,500,707; Beaucage *et al.*, *Tetrahedron Lett.* 22:1859-1862 (1981); Matteucci *et al.*, *J. Amer. Chem. Soc.* 103:3185-3191 (1981); Caruthers *et al.*, *Genetic Engineering* 4:1-17 (1982); Jones, chapter 2, Atkinson *et al.*, chapter 3, and Sproat *et al.*, chapter 4, in Gait, ed. *Oligonucleotide Synthesis: A Practical Approach*, IRL Press, Washington D.C. (1984); Froehler *et al.*, *Tetrahedron Lett.* 27:469-472 (1986); Froehler *et al.*, *Nucleic Acids Res.* 14:5399-5407 (1986); Sinha *et al.*, *Tetrahedron Lett.* 24:5843-5846 (1983); and Sinha *et al.*, *Nucl. Acids Res.* 12:4539-4557 (1984).

Generally, the timing of delivery and concentration of reagents utilized in a coupling cycle will not differ from the protocols typical for unmodified commercial

phosphoramidites used in commercial DNA synthesizers. In these cases, one can merely add the solution containing the pteridine derivatives of this invention to a receptacle on a port provided for an extra phosphoramidite on a commercial synthesizer (*e.g.*, model 380B, Applied Biosystems, Foster City, California, USA). However, where the coupling efficiency of a particular derivatized pteridine compound is substantially lower than the other phosphoramidites, it may be necessary to alter the timing of delivery or the concentration of the reagent in order to optimize the synthesis. Means of optimizing oligonucleotide synthesis protocols to correct for low coupling efficiencies are well known to those of skill in the art. Generally, one merely increases the concentration of the reagent or the amount of the reagent delivered to achieve a higher coupling efficiency. Methods of determining coupling efficiency are also well known. For example, where the 5'-hydroxyl protecting group is a dimethoxytrityl (DMT), coupling efficiency may be determined by measuring the DMT cation concentration in the acid step (which removes the DMT group). DMT cation concentration is usually determined by spectrophotometrically monitoring the acid wash. The acid/DMT solution is a bright orange color. Alternatively, since capping prevents further extension of an oligonucleotide where coupling has failed, coupling efficiency may be estimated by comparing the ratio of truncated to full length oligonucleotide utilizing, for example, capillary electrophoresis or HPLC.

Solid phase oligonucleotide synthesis can be performed using a number of solid supports. A suitable support is one which provides a functional group for the attachment of a protected monomer which will become the 3' terminal base in the synthesized oligonucleotide. The support must be inert to the reagents utilized in the particular synthesis chemistry. Suitable supports are well known to those of skill in the art. Solid support materials include, but are not limited, to polacryloylmorpholide, silica, controlled pore glass (CPG), polystyrene, polystyrene/latex, and carboxyl-modified Teflon. Preferred supports are amino-functionalized controlled pore glass and carboxyl-functionalized Teflon.

Solid phase oligonucleotide synthesis requires, as a starting point, a fully protected monomer (*e.g.*, a protected nucleoside) coupled to the solid support. This coupling is typically through the 3'-hydroxyl (oxo when coupled) covalently bound to a linker which is, in turn, covalently bound to the solid support. The first synthesis cycle then couples a nucleotide monomer, via its 3'-phosphate, to the 5'-hydroxyl of the bound nucleoside through a condensation reaction that forms a 3'-5' phosphodiester linkage

Subsequent synthesis cycles add nucleotide monomers to the 5'-hydroxyl of the last bound nucleotide. In this manner an oligonucleotide is synthesized in a 3' to 5' direction producing a "growing" oligonucleotide with its 3' terminus attached to the solid support.

Numerous means of linking nucleoside monomers to a solid support are known to those of skill in the art, although monomers covalently linked through a succinate or hemisuccinate to controlled pore glass are generally preferred. Conventional protected nucleosides coupled through a hemisuccinate to controlled pore glass are commercially available from a number of sources (e.g., Glen Research, Sterling, Vermont, USA, Applied Biosystems, Foster City, California, USA, Pharmacia LKB, Piscataway, New Jersey, USA).

Placement of a pteridine nucleotide at the 3' end of an oligonucleotide requires initiating oligonucleotide synthesis with a fully blocked furanosyl pteridine linked to the solid support. In a preferred embodiment, linkage of the pteridine nucleoside is accomplished by first derivatizing the pteridine nucleotide as a hemisuccinate. The hemisuccinate can then be attached to amino-functionalized controlled pore glass in a condensation reaction using mesitylene-2-sulfonyl chloride/1-methyl-1H-imidazole as a condensing agent. Controlled pore glass functionalized with a number of different reactive groups is commercially available (e.g., Sigma Chemical, St. Louis, Missouri, USA). A similar coupling scheme is described by Atkinson *et al.*, chapter 3 in Gait, ed., *Oligonucleotide Synthesis: A Practical Approach*, IRL Press, Washington, D.C. (1984). Triisopropylbenzenesulfonyl chloride, imidazolidines, triazolides or even the tetrazolides can also be used as condensing agents. Dicyclohexylcarbodiimide (DCC) and structural analogs are also suitable linkers. Other linkers and appropriate condensing groups are well known to those of skill in the art.

In a preferred embodiment, this invention relates to pteridine nucleotides in which the 3'-hydroxyl is derivatized as a hemisuccinate which may then be covalently bound to a solid support; more specifically to controlled pore glass. In this aspect, R² of Formula I is an amino group which is mono- or di-substituted with a protecting group, such as dimethylaminomethylenamino, or a p-nitrophenylethoxycarbonyl group, and R⁶ is a hemisuccinate, or a hemisuccinate covalently bound to a solid support

It is important to note that where the exocyclic amines are protected by the p-nitrophenylethoxycarbonyl group, the deprotection reagents may also cleave the ester function of the succinyl spacer linking the 3' terminal nucleoside to the solid support. In this case, the coupling scheme described by Stengele *et al.*, *Tetrahedron Lett.* 18:2549-

2552 (1990), is preferred. In this method, solid supports (dihydroxypropyl-CPG, 500Å and 1400Å, Fluka AG, Switzerland, Catalog Nos: 27754, 27764, 2770) are reacted first with N,N'-carbonyldiimidazole and then with 1,6-bismethylaminoheptane as an aliphatic secondary amine spacer. This compound is then coupled with the appropriately protected 2'-nucleoside-3'-O-succinates and the free hydroxyl groups of the solid support are subsequently protected with acetic anhydride and 4-dimethylaminopyridine (DMAP). This linker is completely stable under the deprotection conditions used for p-nitrophenylethoxycarbonyl and p-nitrophenylethyl groups, while cleavage from the matrix can be achieved normally under hydrolytic conditions in concentrated ammonia in less than two hours.

Once the full length oligonucleotide is synthesized, the protecting groups are removed (the oligonucleotide is deprotected), and the oligonucleotide is then cleaved from the solid support prior to use. Where a non-cleavable linker is used, the oligonucleotide can be deprotected and left permanently attached to the support to produce an affinity column. Non-cleavable linkers are well known to those of skill in the art. (See, e.g., WO/ 8501051, EP-A-9174879 and Duncan *et al.*, *Anal. Biochem.*, 169, 104-108 (1988)).

Cleavage and deprotection can occur simultaneously, or sequentially in any order. The two procedures can be interspersed so that some protecting groups are removed from the oligonucleotide before it is cleaved off the solid support and other groups are deprotected from the cleaved oligonucleotide in solution. The sequence of events depends on the particular blocking groups present, the particular linkage to a solid support, and the preferences of the individuals performing the synthesis. Where deprotection precedes cleavage, the protecting groups can be washed away from the oligonucleotide which remains bound on the solid support. Conversely, where deprotection follows cleavage, the removed protecting groups will remain in solution with the oligonucleotide. Often the oligonucleotide will require isolation from these protecting groups prior to use.

In a preferred embodiment, the protecting group on the 5'-hydroxyl is removed at the last stage of synthesis. The oligonucleotide is then cleaved off the solid support, and the remaining deprotection occurs in solution. Removal of the 5'-hydroxyl protecting group typically just requires treatment with the same reagent utilized throughout the synthesis to remove the terminal 5'-hydroxyl groups prior to coupling the next nucleotide monomer. Where the 5'-hydroxyl protecting group is a dimethoxytrityl

group, deprotection can be accomplished by treatment with acetic acid, dichloroacetic acid or trichloroacetic acid.

Typically, both cleavage and deprotection of the exocyclic amines are effected by first exposing the oligonucleotide attached to a solid phase support (via a base-labile bond) to the cleavage reagent for about 1-2 hours. In this manner, the oligonucleotide is released from the solid support. The solution containing the released oligonucleotide is then heated for at least 20-60 minutes at about 80-90°C so that the protecting groups attached to the exocyclic amines are removed. The deprotection step can, alternatively, take place at a lower temperature, but must be carried out for a longer period of time (*e.g.*, the heating can be at 55°C for 5 hours). In general, the preferred cleavage and deprotection reagent is concentrated ammonia.

Where the oligonucleotide is a ribonucleotide and the 2'-hydroxyl group is blocked with a tert-butyldimethylsilyl (TBDMS) moiety, the latter group may be removed using tetrabutylammonium fluoride in tetrahydrofuran at the end of synthesis. *See, Wu et al., J. Org. Chem.* 55:4717-4724 (1990). Phenoxyacetyl protecting groups can be removed with anhydrous ammonia in alcohol (under these conditions the TBDMS groups are stable and the oligonucleotide is not cleaved). The benzoyl protecting group of cytidine is also removed with anhydrous ammonia in alcohol.

Where the exocyclic amines are protected by the p-nitrophenyl-ethoxy-carbonyl group and the coupling to the solid support is via a 1,6-bis-methylamino-hexane condensed with succinate nucleoside, the amino groups are preferably deprotected by treatment with a 1 M DBU (1,8-diaza-bicyclo[5.4.0]-undec-7-ene). Cleavage of the oligonucleotide from the solid support is then accomplished by treatment with concentrated ammonia.

The oligonucleotide of the present invention are not limited to short single stranded sequences. One of skill would recognize that while oligonucleotide synthesis typically has an upper limit of approximately 200 to 500 bases, a number of oligonucleotide can be ligated together to form longer sequences. In addition, oligonucleotide having complementary sequences can be hybridized together to form double-stranded molecules. Methods of hybridizing and ligating oligonucleotide to form longer double stranded molecules are well known (*see, for example, Sambrook et al., Molecular Cloning - A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1985*).

In a preferred embodiment, the label oligonucleotide of this invention double as primers in an amplification reaction discussed in detail below. In this embodiment, one or more amplification primers are provided that each contain at least one compound of Formula I. In a particularly preferred embodiment, the compound of Formula I is one that will permit a polymerase to read through, but not add a corresponding nucleotide in the nucleic acid being extended by the polymerase. In still another embodiment, the label oligonucleotide doubles as an amplification primer and the loop that dequenches the compound of Formula I.

V. *Use of Compounds of Formula I as Fluorescent Labels*

One of skill will recognize that the pteridine derivatives of this invention can be used simply as fluorescent labels to label almost any biological molecule. The unprotected pteridines alone may be linked by the pteridine 1N or 8N, either directly or through a linker or spacer to a composition it is desired to label. Alternatively, the pteridine nucleosides can be used as fluorescent labels. They can be linked preferably through the 5'-hydroxyl, the 3'-phosphate, or the 2'-hydroxyl (in the case of a ribofuranose) directly, or through a linker, to the composition it is desired to label. Such labeled compositions can include, but are not limited to, biological molecules such as antibodies, ligands, cell surface receptors, and enzymes

The nucleic acid can be covalently coupled to the biological molecules either directly via an activated group (*e.g.*, a hydroxyl, a carboxyl) or through a linker that provides reactive moieties that bind to the oligonucleotide and to the biological molecule respectively. Linkers suitable for attaching nucleic acids to biological molecules are well known. Generally, linkers are either hetero- or homo-bifunctional molecules that contain two or more reactive sites that may each form a covalent bond with the respective binding partner (the biological molecule or the nucleic acid). For example, compounds of Formula I can be joined by a peptide linker, by a straight or branched chain carbon chain linker, or by other linkers known by those of skill in the art. Heterobifunctional cross linking reagents such as active esters of N-ethylmaleimide have been widely used. *See, for example, Lerner et al. Proc. Nat. Acad. Sci. (USA), 78:3403-3407 (1981) and Kitagawa et al. J. Biochem., 79:233-236 (1976).* Other linkers, such as those used in the synthesis of nucleic acids are also suitable (*see, e.g., PCT Publication WO 85/01051, Pochet et al. Tetrahedron. 43: 3481-3490 (1987), Schwyzer et al., Helv. Chim. Acta,*

67:1316-1327 (1984), Gait, ed. *Oligonucleotide Synthesis: a Practical Approach*, IRL Press, Washington D.C. (1984)).

VI. *Use of Compounds of Formula I in DNA Amplification*

The compounds of Formula I can be derivitized as nucleotide

5 triphosphates. The nucleotide triphosphate compounds of the present invention can be utilized as monomers for DNA synthesis in DNA amplification techniques such as polymerase chain reaction (*see*, Innis, *et al.*, *PCR Protocols. A Guide to Methods and Application*, Academic Press, Inc., San Diego (1990)), ligase chain reaction (LCR) (*see*, Wu *et al.*, *Genomics* 4:560 (1989), Landegren, *et al.*, *Science* 241:1077 (1988) and
10 Barringer, *et al.*, *Gene* 89:117 (1990)), transcription amplification (*see*, Kwoh *et al.*, *Proc. Natl. Acad. Sci. USA* 86:1173 (1989)) and self-sustained sequence replication (*see*, Guatelli, *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:1874 (1990)). Amplification utilizing the pteridine nucleotides of this invention provides a rapid assay for a particular DNA sequence. Where the presence or absence of a particular DNA sequence is diagnostic of a
15 pathological condition (*e.g.*, AIDS), amplification using the pteridine nucleotide triphosphates provides an extremely sensitive and rapid diagnostic tool.

For example, if PCR amplification is used, a pair of PCR primers will be chosen that are complementary to the DNA sequences flanking the DNA sequence of interest. The PCR reaction mixture will contain one or more species of nucleotide
20 triphosphates of this invention. If the proper target sequences are present in the sample, the DNA sequence between the primers will be amplified. This amplified DNA sequence will contain the pteridine nucleotide triphosphates. The amplified sequence may be separated from the remaining monomers in the mixture by simple size fractionation (*e.g.*, by using an NAP column, Pharmacia-LKB, Piscataway, New Jersey, USA) gel
25 electrophoresis, or other techniques well known to those of skill in the art. The presence or absence of the amplified sequence can then be immediately detected by measuring the fluorescence of the sequence.

Alternatively, fluorescence polarization (FP) measurements can be used to detect a positive or negative PCR reaction without the necessity of separating the PCR
30 products from the primers and nucleotide monomers. The technique uses pteridine nucleotide monomers or, alternatively, relatively short primers, about 25 base pairs each, that incorporate pteridine nucleotide monomers. After the PCR procedure is completed, the resulting mixture is analyzed using FP, by passing a beam of polarized light at an

excitatory wavelength through the mixture. If the target sequence is not present in the starting mixture, the fluorescent primers will remain in solution as relatively small single-stranded fragments, or the fluorescent nucleotide monomers will remain in solution as relatively small molecules. Both the monomers and the short primer fragments will emit a relatively scattered and non-polarized fluorescent light. By contrast, if the target sequence is present, the pteridine monomers or the fluorescent primers will be incorporated into larger double-stranded segments which will move more slowly in response to the excitatory signal and the fluorescent light emitted by the mixture will be more polarized. *See*, EP No. 382433, which describes this technique in greater detail.

Certain nucleotide triphosphate compounds of Formula I are preferred. For instance, compounds of Formula I wherein R^1 is hydrogen or methyl, R^2 is NH_2 , R^3 is methyl, R^7 is a triphosphate, R^5 is hydrogen, and R^6 is hydrogen are preferred.

VII. *Use as Fluorescent Probes*

In another embodiment, this invention provides methods of detecting the presence, absence, or quantity of a target nucleic acid by utilizing a nucleic acid that contains one or more compounds of Formula I. The fluorescence of a compound of Formula I is quenched when it is incorporated into an oligonucleotide (*see, e.g.*, U.S. Patent No. 5,525,711 and Hawkins *et al.* (1995) *Nucl. Acids Res.* 23:2872-2880). However, when the compound of Formula I is present as an insertion, fluorescence activity is partially or completely restored. Without being bound by a particular theory, it is believed that alteration of the normal conformation (*e.g.*, base stacking) of the oligonucleotide at the location of the compound of Formula I reduces and/or eliminates the quench thereby causing an increase in fluorescence.

The methods involve contacting the target nucleic acid with a nucleic acid probe where the nucleic acid probe comprises a compound of Formula I located in the probe such that, when the probe hybridizes to the target nucleic acid, the compound of Formula I is present in a loop that does not participate in complementary base pairing with a nucleotide of the target nucleic acid. Next, detecting the fluorescence produced by the compound of Formula I when the probe forms a hybrid duplex with the target nucleic acid.

In certain embodiments, the nucleotide sequences of the oligonucleotide of this invention can be selected such that the normal base stacking of the compound of Formula I and/or its adjacent nucleotide is disrupted either through self-hybridization of

the oligonucleotide or when the oligonucleotide hybridizes to a target nucleic acid or when the oligonucleotide participates in a protein/DNA interaction. This is typically accomplished when the compounds of Formula I are located in a loop(s).

5 The term nucleic acid loop, as used herein, refers to a region of one or more contiguous nucleic acids that do not participate in normal (*e.g.*, purine/pyrimidine hydrogen bond formation) base pairing when the nucleic acid is hybridized to an otherwise complementary target nucleic acid. In a preferred embodiment, the loops of this invention are formed as insertions in nucleic acid sequences. The insertion can comprise one or more nucleotides. As used herein an insertion refers to the addition of
10 one or more nucleotides into an oligonucleotide that is otherwise complementary to a target nucleic acid or target nucleotide subsequence. The insertion is thus recognized by reference to the target nucleic acid sequence or subsequence. One of skill will appreciate that an insertion need not be produced by actual physical insertion of one or more additional nucleotides (bases) into an already existing oligonucleotide, but reflects the presence of the extra nucleotide(s) with reference to a particular target sequence or
15 subsequence. Thus, the insertion-containing oligonucleotide can be synthesized *de novo*. Alternatively, the insertion can be created by deleting one or more nucleotides in the target sequence or subsequence, or by synthesizing (*e.g.*, polymerizing) a target sequence or subsequence lacking nucleotides corresponding to one or more nucleotides in the label
20 oligonucleotide.

In one preferred embodiment, the loop ranges in length from about 1 to about 100 nucleotides when the probe hybridizes to the target nucleic acid. In particularly preferred probes, the loop is an insertion in the nucleic acid probe which is otherwise complementary to the target nucleic acid or to a contiguous subsequence of the
25 target nucleic acid. In some preferred embodiments, the insertion is three nucleotides in length and comprises two nucleotides each adjacent to the compound of Formula I. In particularly preferred embodiments, at least one nucleotide adjacent to the compound of Formula I is a purine (*e.g.*, adenosine), or is bordered by at least two adjacent purines (*e.g.*, adenosine) in both the 5' and 3' direction. In another preferred embodiment, the
30 insertion is a single base insertion, the compound of Formula I.

In yet another embodiment, the insertion is self-complementary and forms a hairpin in which the compound of Formula I is present in the loop of the hairpin and does not participate in complementary base pairing. The nucleotides comprising the loop can be selected such that they are not complementary to the corresponding nucleotides of

the target nucleic acid when the probe is hybridized to the target nucleic acid and where the probe is complementary to at least two non-contiguous subsequences of the target nucleic acid.

The hairpin conformation can be one that is stabilized by hybridization to the target nucleic acid, or it can be a stable formation in the label oligonucleotide even when the label oligonucleotide is not hybridized to the target substrate. In this case, the base stacking of the label oligonucleotide is always disrupted resulting in a continually elevated fluorescence activity. The label oligonucleotide is an "always-on" label. The hairpin formation can be internal to the label oligonucleotide or alternatively can be formed at the terminus of the label oligonucleotide. The label oligonucleotide of this invention can include one or more of the above-described loops. In addition, any loop can include one or more compounds of Formula I. In another embodiment, the compound of Formula I is present in a terminal subsequence of the nucleic acid probe where the terminal subsequence does not hybridize to the target nucleic acid when the remainder of the nucleic acid probe hybridizes to the target nucleic acid. The terminal subsequence can form a terminal hairpin by hybridization with a second subsequence of the probe such that the compound of Formula I is present in a loop of the hairpin and does not participate in complementary base pairing.

One of skill in the art will appreciate that the methods and molecules of this invention can be used in a wide variety of contexts. Under conditions that permit self-hybridization, normal base stacking of compounds of Formula I is disrupted and the nucleotide fluoresces. These oligonucleotide continuously fluoresce and provide useful "always on" labels. Always on labels can also be produced by placing the compound of Formula I at the 3' or 5' terminus, most preferably at the 5' terminus. In this position, the compound of Formula I is not as quenched as when it is located within the oligonucleotide. The "always on" fluorescent labels can be used in a manner analogous to fluorescent labels (*e.g.*, fluorescein, rhodamine, *etc.*) known in the prior art.

In a more preferred embodiment, the compounds of this invention are used to label molecules, more preferably biological molecules such as other nucleic acids, proteins, and the like. Particularly preferred molecules to be labeled include antibodies, growth factors, cell-surface receptors, lectins, hormones, and the like. Indeed, the molecules that can be so labeled include virtually any molecule that can be linked to a nucleic acid. The oligonucleotide of this invention can be linked to the subject molecules, *e.g.*, proteins, or nucleic acids, directly, or through a linker.

Suitable linkers attaching nucleic acids to other molecules are well known to those of skill in the art. Generally linkers are either hetero or homobifunctional molecules that contain two or more reactive sites that can form a covalent bond with the nucleic acid and the molecule to which it is to be attached. For example, the label nucleic acids of this invention can be joined to the subject molecule by a peptide linker, by a straight or branched chain carbon chain linker, or by a heterocyclic carbon. The linkers can attach to convenient reactive moieties on the base (*e.g.*, NH_3) to available hydroxyl groups on the ribose or to a terminal phosphate. Heterobifunctional cross linking reagents such as active esters of ethylmaleimide have been widely used (*see, e.g.*, Lerner *et al.* (1981) *Proc. Nat. Acad. Sci. USA* 78:34033407 and Kitagawa *et al.* (1976) *J. Biochem.* 79:233236), and other suitable linkers are well known to those of skill in the art (*see, e.g.*, Chapter 4 In *Monoclonal Antibodies, Principles and Applications*, Birch and Lennox, eds., Wiley-Liss, Inc., New York (1994)).

In another embodiment, the oligonucleotide of this invention have a nucleotide sequence that enables the molecules to act as molecular beacons. The term "molecular beacon," as used herein refers to a molecule capable of participating in a specific binding reaction and whose fluorescence activity changes when the molecule participates in that binding reaction. Preferred molecular beacons of this invention are label oligonucleotide that show little or no fluorescence activity when the molecule is free in solution and yet show a detectable increase in fluorescence activity when bound to their target substrate. The molecules are designed so that interaction (*e.g.*, binding) with the substrate (target) molecule introduces a change in conformation of the label oligonucleotide that results in a reduction of the quenching of the compound of Formula I present in the label oligonucleotide. Such changes in conformation preferably involve a disruption of the normal base stacking of the compound of Formula I in the label oligonucleotide. Such disruptions are preferably produced by a one or more base pair mismatch between the label oligonucleotide and its target when the molecules are hybridized, by the formation of loops (*e.g.*, hairpins) in the label oligonucleotide, by lack of complementarity between the compound of Formula I and the corresponding nucleotides in the target

The compounds of Formula I possess fluorescence quantum yields higher than other structurally similar pteridine nucleotides.

Particularly preferred molecular beacons have a relative quantum yield in the unbound (quenched) state that ranges from undetectable to about 0.8, more preferably from undetectable to about 0.1 M and most preferably from undetectable to about 0.05.

Virtually any detectable change in fluorescence on binding is useful.

- 5 However, the larger the change in fluorescence, the easier it is to detect the binding event. Thus, preferred molecular beacons show a 2-fold increase, more preferably at least a 5-fold increase and most preferably at least about 10 to about 20-fold increase in fluorescence intensity on binding to the target molecule.

10 The label oligonucleotide can also be used to detect interactions with nucleic acid binding proteins (or other molecules). The compounds of Formula I present in the label oligonucleotide are located so that the normal planar base stacking is disrupted when the label oligonucleotide is bound by a protein (*e.g.*, rec A protein, PI nuclease, HIV integrase, estrogen receptor, etc.). Again, the disruption reduces or eliminates the "quench" resulting in an increase in fluorescence activity of the
15 protein/label oligonucleotide complex. The increased fluorescence can be easily detected as discussed below. The label oligonucleotide sequence is selected so that the label oligonucleotide is recognized and bound by the particular protein (or other molecule) of interest.

20 In another preferred embodiment, the label oligonucleotides of this invention provide a means of detecting the presence and/or absence and/or quantifying the product of a nucleic acid amplification reaction. In this embodiment, the nucleotide sequence of the label oligonucleotide is selected so that the base stacking of the compound of Formula I is not disrupted unless an amplification product is present.

25 In one embodiment, the oligonucleotide does not participate in the amplification reaction. It is simply present as a separate indicator molecule. In this context, the label oligonucleotide nucleotide sequence is selected so that a loop containing the compound of Formula I is formed when the label oligonucleotide hybridizes to the amplification product. As amplification product is formed, the label oligonucleotide hybridizes to that product, disrupting the base pair stacking and thereby
30 increasing its fluorescence activity. As long as the label oligonucleotide is present in a molar excess (of the amplification product) the change in fluorescence intensity in the reaction vessel is proportional to the amount of amplification product. However, it is also desirable to maximize the signal to noise ratio. Therefore, in a preferred embodiment, the

molar excess of label oligonucleotide is kept as low as possible to minimize generation of a background signal.

In another embodiment, the compound of Formula I can be incorporated into one or more of the amplification primers such that when the primer is extended by formation of an amplification product, the extended primer forms a loop thereby dequenching the compound of Formula I.

In other embodiments, the fluorescent labels of this invention are particularly well suited for the detection of nucleic acid hybridization in a wide variety of contexts including, but not limited to, nucleic acid hybridization arrays, Southern blot hybridizations, *in situ* hybridization, and the like.

One of skill will appreciate that the fluorescent probes of the present invention can be immobilized on a substrate. The substrates include, but are not limited to, a glass substrate, (e.g., a high density array), a solid support, or a gel. A high density array can be synthesized on a substrate by attaching photoremovable groups to the surface of a substrate, exposing selected regions of the substrate to light to activate those regions, attaching a nucleic acid monomer (e.g., a compound of Formula I) with a photoremovable group to the activated regions, and repeating the steps of activation and attachment until probes of the desired length and sequences are synthesized. The resulting array of probes can then be used to determine target nucleic acids. (See, e.g., Fodor *et al. Science*, 251:767-773 (1991) and U.S. Patent No. 5,143,854.).

VIII. Hybridization with Label Oligonucleotides.

The label oligonucleotide of this invention can be used as "always on" labels and simply attached to the molecule or article of interest just like any other fluorescent marker. Alternatively, the label oligonucleotides are used as molecular beacons whose fluorescence activity increases when the molecules hybridize to a target nucleic acid or nucleic acid subsequence or when the label oligonucleotides are bound by a target protein.

The nucleic acid hybridization simply involves providing a denatured label oligonucleotide probe and target nucleic acid under conditions where the probe and its target can form stable hybrid duplexes through complementary base pairing. As described above, the label oligonucleotide have nucleic acid sequences that introduce a change in conformation on hybridizing that increases the fluorescence activity of the

compound of Formula I present in the label oligonucleotide. The resulting fluorescent hybrid duplexes are then detected, *e.g.*, using a spectrofluorometer.

It is generally recognized that nucleic acids are denatured by increasing the temperature or decreasing the salt concentration of the buffer containing the nucleic acids.

5 Under low stringency conditions (*e.g.*, low temperature and/or high salt) hybrid duplexes (*e.g.*, DNA:DNA, RNA:RNA, or RNA:DNA) will form even where the annealed sequences are not perfectly complementary. Thus specificity of hybridization is reduced at lower stringency. Conversely, at higher stringency (*e.g.*, higher temperature or lower salt) successful hybridization requires fewer mismatches.

10 One of skill in the art will appreciate that hybridization conditions can be selected to provide any degree of stringency. In a preferred embodiment, hybridization is performed at low stringency (*e.g.*, about 20°C to about 50°C, more preferably about 30°C to about 40°C, and most preferably about 37°C and 6X SSPE-T or lower for an oligonucleotide) to ensure hybridization and then subsequent washes are performed at
15 higher stringency (*e.g.*, typically, stringent conditions will be those in which the salt concentration is at most about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (*e.g.*, 10 to 50 nucleotides) and at least about 50°C or 60°C for longer probes). Successive hybridizations can be performed at increasingly higher stringency (*e.g.*, down to as low as
20 0.25 X SSPE-T at 37°C to 50°C) until a desired level of hybridization specificity is obtained. Stringency can also be increased by addition of agents such as formamide.

Hybridization specificity can be evaluated by specific labeling of nucleic acids separated in gel electrophoresis and/or by evaluation of the signal to noise (background fluorescence), or by other methods well known to those of skill in the art. In
25 general, there is a tradeoff between hybridization specificity (stringency) and signal intensity. Thus, in a preferred embodiment, the wash is performed at the highest stringency that produces consistent results and that provides a hybridization showing a readily detectable change in signal intensity (*e.g.*, an increase in signal intensity of at least 20%, more preferably an increase in signal intensity of at least 50% and most preferably
30 an increase in signal intensity of at least 100% (a doubling of signal intensity) over the background signal intensity.

Methods of selecting and optimizing nucleic acid sequences for hybridization to particular targets and/or internal hybridization and/or priming of particular templates are well known to those of skill in the art (*see, e.g.*, Innis *et al.* (1990)

PCR Protocols, A Guide to Methods and Applications, Innis *et al.*, Academic Press, Inc., N.Y.). Such optimization is simplified by the use of nucleic acid sequence analysis software. Such software is well known to those of skill in the art and includes, but is not limited to, HyperPCR, Loop Viewer, MulFold, Primer, and Amplify (available on the internet at PubNet), GeneWorks7, GeneJockey, DNA Strider, LaserGene (DNAS⁺Star, Madison, Wisconsin, USA), and the like.

IX. *Signal Detection.*

Means of detecting the fluorescent signals produced by the label oligonucleotide of this invention are well known to those of skill in the art. Labeled target (sample) nucleic acids hybridized to the probes of the high density array are known to those of skill in the art. Typically detection is the same as that for any other fluorescent label. Such detection involves exposing the fluorescent moiety (*i.e.*, the label oligonucleotide) to an excitation illumination at the absorption wavelength of the compound of Formula I. The light is absorbed and re-emitted at the emission wavelength of the compound of Formula I. Devices for detecting fluorescent labels are commercially available and include, but are not limited to, fluorometers, fluorescence microscopes, flow cytometers, fluorescence plate readers, and the like (*see, e.g.*, Applied Imaging Corp., Santa Clara, California, USA; Perkin-Elmer Corp., Norwalk, Connecticut, USA; and Photon Technology International, South Brunswick, New Jersey, USA).

In a particularly preferred embodiment, fluorescence is detected using a PTI (Photon Technologies, Inc., New Brunswick, New Jersey, USA) spectrofluorometer with a double excitation monochromater, a water cooled photomultiplier, and a sample chamber coupled with a water bath (Pharmacia LKB, Piscataway, New Jersey, USA). Low volume measurements can be made using an "H" style cuvette with excitation over the long (*e.g.*, 1 cm) path and emission through the short (*e.g.*, 2 mm path). In another embodiment, 3 mm x 3 mm square cuvettes with a brass adaptor have been successfully used.

In another embodiment the fluorescence of the label oligonucleotide can be detected indirectly through energy transfer through a second and even a third fluorophore. In this embodiment, a second fluorophore is provided that has an absorption wavelength at or about the emission wavelength of the compound of Formula I. When compound of Formula I is excited, the energy it releases is absorbed (*e.g.*, through resonance energy transfer) by the second fluorophore which then fluoresces at its

characteristic wavelength. This approach is particularly convenient where it is desired to shift the signal to a wavelength different than the characteristic fluorescence wavelength of the compound of Formula I. Resonance energy transfer systems are well known to those of skill in the art (*see, e.g.*, Forster (1948) *Ann. Phys.* 2:55; Stryer *et al.* (1967)

5 *Proc. Natl. Acad. Sci. USA* 58:719-726, and Stryer (1978) *Ann. Rev. Biochem.* 47:819).

X. Kits

10 An additional aspect of the invention relates to kits useful for the detection of nucleic acid/nucleic acid interactions and/or for the detection of protein/nucleic acid interactions and/or for the detection of amplification product. These kits take a variety of forms and can comprise one or more containers containing one or more label oligonucleotide of this invention. The label oligonucleotide can be "always on" probes or "molecular beacons" as described above. The label oligonucleotide can be simple "indicators" of a particular nucleic acid or protein target, or they can be amplification primers for amplification and detection of a particular target nucleic acid. Other optional components of the kit include, for example, a polymerase, a reaction vessel, a second fluorophore for detection through resonance energy transfer, the appropriate buffers for PCR or other amplification reactions, positive and negative controls for diagnostic application, and the like. In addition to the above components, the kit can also contain instructions for use.

15 In other aspects of this invention, the kits can be used for the synthesis of oligonucleotides. In this aspect, the kit comprises compounds of Formula I, preferably wherein R⁶ is a phosphoramidite derivative.

20 In other aspects, the kits can be used to generate fluorescent labels. In this embodiment, the kit comprises a compound of Formula I and optionally contains a linked compound. Such linked compounds include, but are not limited to, biological molecules such as antibodies, lipids, liposomes, ligands, polysaccharides, cell surface receptors, and enzymes. These and other uses for kits will be readily apparent to those of skill in the art.

EXAMPLES

Example 1

This example illustrates the synthesis of the pteridone bases.

A. 4-Amino-6-methyl-7(8H)-pteridone (3). [D. Söll *et al.*, *Chem. Ber.*

96:2977 (1963)]

4,5,6-Triaminopyrimidine (1) [J. Baddiley *et al.*, *J. Chem. Soc.* 386 (1943)] (2.0 g, 16 mmoles) and ethyl pyruvate (2.2 mL, 19 mmoles) were heated in glacial acetic acid (20 mL) for 2 hours. After cooling the precipitate was collected, washed with water and purified by recrystallization from DMF/ H₂O (350 mL, 1:1). Yield: 1.25 g (44%). M.p. >360°C. UV (pH 5), λ_{max} (log ϵ): 218 (4.37), 243 (4.07), [291 (3.82)], 326 (4.02), [343 (3.89)]. ¹H-NMR (D₆-DMSO): 2.34 (s, MeC(6)); 7.35 + 7.45 (2 bs, NH₂); 8.12 (s, H-C(2)) 12.56 (bs, H-N(8)).

B. 4-Amino-2,6-dimethyl-7(8H)-pteridone (4).

4,5,6-Triamino-2-methylpyrimidine (2) [B. Lythgoe *et al.*, *J. Chem. Soc.* 315 (1944)] (0.5 g, 3.6 mmoles) and ethyl pyruvate (1 mL) were heated in AcOH (10 mL) and EtOH (10 mL) under reflux for 1 hour. After cooling the precipitate was collected and recrystallized from DMF/ H₂O (90 mL, 1:1) with a little charcoal to give 0.372 g (54%) of colorless crystals. M.p. >300°C (decomp.). UV (pH 5), 245 (4.08), [291 (3.77)], [318 (4.00)], 331 (4.03), [345 (3.88)]. ¹H-NMR (D₆-DMSO): 2.34 (s, 6H, Me-C(2), Me-C(6)), 7.32 + 7.42 (2 bs, NH₂), 12.43 (bs, H-N(8)). Anal. calcd. for C₈H₉N₅O (190.1): C 50.26, H 4.74, N 36.63; found: C 50.15, H 4.82, N 36.35.

Example 2

This example illustrates the coupling of a pteridone base to a deoxyribose.

A. 4-Amino-6-methyl-8(2-deoxy-3,5-di-O-p-chlorobenzoyl- β -D-ribofuranosyl)-7(8H)-pteridone (5).

4-Amino-6-methyl-7(8H)-pteridone (3) (1.54 g, 9 mmoles) and DBU (1.34 mL, 9 mmoles) were stirred in anhydrous acetonitrile (100 mL) for 30 min. Then 2-deoxy-3,5-di-O-p-chlorobenzoyl- α -D-ribofuranosyl chloride [J. J. Fox *et al.*, *J. Am. Chem. Soc.* 83:4066 (1961)] was added and stirring continued for 2 hours. It was evaporated, the residue dissolved in dichloromethane (60 mL), washed with a saturated solution of NaCl (2 x 30 mL), the organic layer dried over Na₂SO₄, filtered and again evaporated. The crude product was dissolved in toluene (10 mL), put onto a silica gel

column (5 x 11 cm) and eluted with toluene/EtOAc (2:1, 600 mL) and toluene/EtOAc (1:1, 400 mL). The main fraction was evaporated and then recrystallized from CHCl₃/MeOH to give 1.7 g (34%) colorless crystals. M.p. >187-189°C. UV (MeOH): 241 (4.66), 333 (3.96). ¹H-NMR (D₆-DMSO): 2.38 (s, Me-C(6)), 2.55 (m, H_α-C(2')), 3.18 (m, H_β-C(2')), 4.53 (m, 2 H-C(5')), 4.69 (m, H-C(4')), 5.94 (m, H-C(3')), 7.34 (m, H-C(1')), 7.51 (d, 2H p-Clbz), 7.59 (d, 2H p-Cl-bz), 7.79 (bs, NH₂), 7.93 (2d, 4H p-Clbz), 8.22 (s, H-C(2')).

Anal. calcd. for C₂₆H₂₁Cl₂N₅O₆ (570.4): C 54.75, H 3.71, N 12.28; found: C 54.52, H 3.80, N 12.28.

10 B. *4-Amino-2,6-dimethyl-8-(2-deoxy-3,5-di-O-p-chlorobenzoyl-β-D-ribofuranosyl)-7(8H)-pteridone (6).*

4-Amino-2,6-dimethyl-7(8H)-pteridone (4) (1.58 g, 8.26 mmoles) was suspended in anhydrous acetonitrile (60 mL), DBU (1.23 mL, 8.26 mmoles) added and stirred for 15 min at room temperature. Then 2-deoxy-3,5-di-O-p-chlorobenzoyl-α-D-ribofuranosyl chloride (4.2 g, 10 mmoles) was added and the mixture stirred for 2 hours. The colorless precipitate was collected and recrystallized from CHCl₃/MeOH (1:2, 90 mL) to give 1.92 g (40%). M.p. >218-220°C (decomp). UV (MeOH): 241 (4.66), 333 (3.96). ¹H-NMR (D₆-DMSO): 2.36 (s, Me-C), 2.40 (s, MeC), 2.57 (m, H_α-C(2')), 3.17 (m, H_β-C(2')), 4.53 (m, 2 H-C(5')), 4.69 (m, H-C(4')), 6.00 (m, H-C(3')), 7.30 (m, H-C(1')), 7.42 (bs, NH₂), 7.50 (d, 2H p-Clbz), 7.59 (d, 2h p-Clbz), 7.89 (d, m2H p-Clbz), 7.96 (d, 2H p-Clbz).

Anal. calcd. for C₂₇H₂₃Cl₂N₅O₆ (548.4): C 55.49, H 3.97, N 11.98; found: C 55.57, H 4.09, N 11.53.

25 C. *4-Amino-6-methyl-8-(2-deoxy-β-D-ribofuranosyl)-7(8H)-pteridone (7).*

Compound (5) (0.5 g, 0.88 mmoles) was added to a solution of sodium (20 mg) in anhydrous MeOH (20 mL) and stirred at room temperature for 12 hours. It was then neutralized by AcOH and the suspension concentrated to 10 mL. The precipitate was collected, washed with MeOH and dried in high vacuum to give 0.16 g (62%). Cooling of the filtrate provided a second crop 0.05 g (19%). M.p. >190°C decomposition. UV (MeOH): 248 (4.09), [292 (3.71)], 329 (3.93). ¹H-NMR (D₆-DMSO): 2.01 (m, H_α-C(2')), 2.36 (s, Me), 2.88 (m, H_β-C(2')), 3.54 (m, H-C(5')), 3.60 (m, H-C(5')), 3.67 (m, H-C(4')), 4.44 (m, H-C(3')), 4.72 (dd, HO-C(5')), 5.17 (d, HO-C(3')), 7.15 (m, H-C(1')), 7.52 + 7.70 (2s, NH₂), 8.20 (s, H-C(2')).

Anal. calcd. for $C_{12}H_{15}N_5O_4$ (293.3): C 49.13, H 5.16, N 23.88; found: C 49.34, H 5.18, N 23.78.

D. *4-Amino-2,6-dimethyl-8-(2-deoxy- β -D-ribofuranosyl)-7(8H)-pteridone (8).*

Compound (6) (0.78 g, 1.33 mmol) was added to a solution of sodium (30 mg) in anhydrous MeOH (20 mL) and stirred at room temperature for 12 hours. It was then neutralized by AcOH and the suspension concentrated to 10 mL. The precipitate was collected, washed with MeOH and dried in high vacuum to give 0.38 g (93%) of colorless crystals. M.p. >180°C decomposition. UV (MeOH): 250 (4.08), [298 (3.74)], 333 (3.95). 1H -NMR (D_6 -DMSO): 1.99 (m, H_α -C(2')), 2.34 (s, Me), 2.37 (s, Me), 2.88 (m, H_β -C(2')), 3.53 (m, H-C(5')), 3.65 (m, H-C(5'')), 3.74 (m, H-C(4')), 4.45 (m, H-C(3')), 4.66 (dd, HO-C(5')), 5.16 (d, HO-C(3')), 7.14 (m, H-C(1')), 7.38 + 7.57 (2s, NH_2). Anal. calcd. for $C_{13}H_{17}N_5O_4$ (307.3): C 50.81, H 5.58, N 22.79; found: C 50.34, H 5.80, N 23.08.

Example 3

This example illustrates the protection of O-5' of the deoxyribose.

A. *4-Amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-ribofuranosyl]-7(8H)-pteridone (9).*

Compound (7) (1.25 g, 4.26 mmol) was twice coevaporated with anhydrous pyridine (20 mL) and then dissolved in the same solvent (20 mL). Then 4,4'-dimethoxytrityl chloride (1.73 g, 5.1 mmol) was added and stirred at room temperature for 12 hours. It was evaporated, the residue dissolved in CH_2Cl_2 (40 mL) and washed with saturated $NaHCO_3$ solution (2 x 20 mL). The organic phase was dried over Na_2SO_4 , evaporated, the residue dissolved in toluene (5 mL) and put onto a silica gel column for chromatography with toluene/EtOAc (1:3, 400 mL) and toluene/EtOAc (1:1, 600 mL). The product fraction was evaporated and the resulting solid foam dried in high vacuum to give 2.16 g (85%). M.p. >90-105°C. UV (MeOH): [233 (4.52)], [278 (3.84)], 328 (3.94). 1H -NMR (D_6 -MSO): 2.11 (m, H_α -C(2')), 2.30 (s, Me-C(6')), 2.78 (m, H_β -C(2')), 3.15 (m, H-C(5')), 3.33 (m, H-C(5'')), 3.69 (s, OMe), 3.70 (s, OMe), 3.94 (m, H-C(4')), 4.44 (m, H-C(3')), 5.17 (d, OH-C(3')), 6.77 (2d, 4H, trityl), 7.20 (m, 7H, trityl), 7.35 (m, 2H, trityl), 7.50 + 7.67 (2 bs, NH_2), 8.11 (s, H-C(2)).

Anal. calcd. for $C_{33}H_{33}N_5O_6$ (595.7): C 66.54, H 5.58, N 11.76; found: C 66.58, H 5.92, N 11.73.

B. *4-Amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-ribofuranosyl]-7(8H)-pteridone (10).*

5 Compound (8) (0.57 g, 1.85 mmoles) was twice coevaporated with anhydrous pyridine (10 mL) and then dissolved in the same solvent (15 mL). Then 4,4'-dimethoxytrityl chloride (0.752 g, 2.22 mmoles) was added and stirred at room temperature for 12 hours. It was evaporated, the residue dissolved in CH_2Cl_2 (20 mL) and washed with saturated $NaHCO_3$ solution (2 x 20 mL). The organic phase was dried
10 over Na_2SO_4 , evaporated, the residue dissolved in toluene (5 mL) and put onto a silica gel column for chromatography with toluene/EtOAc (1:1, 300 mL) and toluene/EtOAc (1:3, 200 mL). The product fraction was evaporated and the resulting solid foam dried in high vacuum to give 0.785 g (70%). M.p. >105-120°C. UV (MeOH): [232 (4.46)], [258 (4.11)], 333 (3.96). 1H -NMR (D_6 -DMSO): 2.08 (m, H_α -C(2')), 2.18 (s, Me-C(6)), 2.29 (s, MeC(6)), 2.76 (m, H_β -C(2')), 3.12 (m, H-C(5')), 3.39 (m, H-C(5'')), 3.68 (s, OMe), 3.70 (s, OMe), 3.94 (m, H-C(4')), 4.45 (m, H-C(3')), 5.16 (d, OH-C(3')), 6.78 (2d, 4H, trityl), 7.17 (m, 7H, trityl), 7.33 (m, 2H, trityl), 7.34 + 7.67 (2 bs, NH_2).
15 Anal. calcd. for $C_{33}H_{33}N_5O_6 \cdot x H_2O$ (627.7): C 65.06, H 5.94, N 11.16; found: C 64.92, H 5.92, N 11.23.

20

Example 4

This example illustrates the protection of O-3 of the deoxyribose.

A. *4-Amino-6-methyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)- β -D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H- β -cyanoethyl N-diisopropyl)phosphoramidite (11).*

25

Compound (9) (1.25 g, 2.1 mmoles), bis-(N,N-diisopropylamino)-H- β -cyanoethoxyphosphane (0.95 g, 3.1 mmoles) and 1H-tetrazole (74 mg., 1.05 mmoles) were dissolved in anhydrous CH_2Cl_2 (30 mL) and stirred at room temperature under
30 nitrogen atmosphere for 12 hours. It was diluted with CH_2Cl_2 (10 mL), washed with a saturated $NaHCO_3$ solution (2 x 10 mL), the organic phase dried over Na_2SO_4 and then evaporated. The solid foam was dissolved in toluene (3 mL), put onto a silica gel column (12 x 5 cm) for chromatography with toluene/EtOAc (1:1, 500 mL, containing a few drops of triethylamine). The product fraction was again evaporated to give 1.46 g (86%)

of a solid colorless foam. M.p. >70-75°C. UV (MeOH): [233 (4.49)], [278 (3.81)], 328 (3.91). ¹H-NMR (D₆-DMSO): 0.92-1.07 (m, 2 CHMe₂), 2.29 (s, Me-C(6)), 2.30 (m, H_α-C(2')), 2.59 (t, OCH₂CH₂CN), 2.85 (m, H_β-C(2')), 3.12 (m, H-C(5')), 3.27 (m, H-C(5'')), 3.38-3.57 (m, 2 CHMe₂, OCH₂CH₂CN), 3.68 (s, 2 OMe), 4.06 (m, H-C(4')), 4.71 (m, H-C(3')), 6.76 (m, 4H, trityl), 7.19 (m, 7H, trityl, H-C(1')), 7.33 (m, 2H, trityl), 7.51 + 7.68 (2 bs, NH₂), 8.09 (s, H-C(2)).

Anal. calcd. for C₄₂H₅₀N₇O₇P (795.9): C 63.38, H 6.33, N 12.32; found: C 63.62, H 6.39, N 11.84.

10 B. 4-Amino-2,6-dimethyl-8-[-2-deoxy-5-O-(4,4'-dimethoxytrityl)-β-D-ribofuranosyl]-7(8H)-pteridone-3'-O-(H-β-cyanoethyl N-diisopropyl)phosphoramidite (12).

Compound (10) (0.61 g, 1 mmol), bis-(N,Ndiisopropylamino)-H-β-cyanoethoxyphosphane (0.45 g 1.5 mmoles) and ¹H-tetrazole (35 mg, 0.5 mmoles) were dissolved in anhydrous CH₂Cl₂ (15 mL) and stirred at room temperature under nitrogen atmosphere of 12 hours. It was diluted with CH₂Cl₂ (10 mL), washed with a saturated NaHCO₃ solution (2 x 10 mL), the organic phase dried over Na₂SO₄ and then evaporated. The solid foam was dissolved in toluene (5 mL), put onto a silica gel column (12 x 1.5 cm) for chromatography with toluene/EtOAc (3:2, 150 mL, containing a few drops of triethylamine). The product fraction was again evaporated to give 0.69 g (85%) of a solid colorless foam. UV (MeOH): [231 (4.48)], [258 (4.13)], 334 (3.95). ¹H-NMR (D₆-DMSO): 0.88-1.01 (m, 2 CHMe₂), 2.29 (s, Me-C(2)), 2.30 (s, Me-C(6)), 2.30 (m, H_α-C(2')), 2.59 (t, OCH₂CH₂CN), 2.83 (m, H_β-C(2')), 3.11 (m, H-C(5')), 3.25 (m, H-C(5'')), 3.38-3.57 (m, 2 CHMe₂, OCH₂CH₂CN), 3.69 (s, 2 OMe), 4.10 (m, H-C(4')), 4.73 (m, H-C(3')), 6.75 (m, 4H, trityl), 7.19 (m, 7H, trityl, H-C(1')), 7.33 (m, 2H, trityl), 7.33 + 7.54 (2 bs, NH₂).

Anal. calcd. for C₄₃H₅₂N₇O₇P (809.9): C 63.77, H 6.47, N 12.11; found: C 63.70, H 6.50, N 12.01.

Example 5

30 This example sets forth the synthesis of oligonucleotides using compounds of Formula I.

Oligonucleotide Synthesis and Purification

Probes were synthesized as phosphoramidites and incorporated directly into oligonucleotides using an Applied Biosystems Model 392 (Foster City, CA) automated DNA synthesizer following standard protocols recommended by the

35

manufacturer. The fluorophore phosphoramidite was placed in bottle position 5 on the synthesizer and treated in the same way as the standard phosphoramidites. Table 2 lists sequences of oligonucleotides made using monomers of the present invention. The probe-containing synthetic oligonucleotides were treated in the same way as standard oligonucleotides throughout synthesis and de-blocking procedures. Purification of all strands was done by 20% denaturing polyacrylamide gel electrophoresis (19:1 acrylamide:bis). Bands were detected by UV shadowing, excised, and extracted using an Elutrap (Schleicher and Schuell, Concord, NH) electrical elution device. Double strands were formed by combining complementary strands at equimolar concentrations, heating to > 85°C and allowing to cool to room temperature. Comparison of fluorescence intensity between single and double stranded oligonucleotides was done after adding an excess of non-fluorophore containing strand to insure that all of the fluorophore containing strands were annealed.

15 Melting Temperatures

Melting temperatures of double-stranded oligonucleotides were measured by monitoring absorbance hyperchromicity at 260 nm in an HP spectrophotometer equipped with a Hewlett-Packard 89090A Peltier temperature controller. Samples were measured in 10 mM Tris, pH 7.5, with a NaCl concentration of 10 mM. Temperature was increased by 1°C per minute with a 1 min equilibration time between measurements.

P1 Nuclease Digestion

P1 Nuclease was from Penicillium citrinum (Boehringer Mannheim Biochemica). Oligonucleotides were digested with P1 nuclease using 3 Units in a total volume of 100 µl. A fluorescence scan of starting material was compared with the fluorescence scan of the products after incubation at 37° for 19 hours. The ratio of these two scans was compared to the ratio of the relative quantum yield of the oligonucleotide used in the starting material and the relative quantum yield of the monomer form of the fluorophore being examined.

30 Spectroscopic Analysis

Fluorescence measurements were done on a PTI (Photon Technologies, Inc., New Brunswick, NJ) spectrofluorometer equipped with a 75 Watt xenon arc lamp

using a double excitation monochromator and a water cooled photomultiplier. The system was interfaced with a Pharmacia Multitemp LKB water bath temperature controller (Pharmacia). Small volume samples were measured in 3 mm x 3 mm quartz cuvettes using custom built brass adaptors. Relative quantum yield measurements were done on samples measuring < 0.12 at the excitation wavelength in 10 mM Tris, pH 7.5, at room temperature.

Excitation was at 350 nm and the emission wavelength was 450 nm. Corrected emission spectra were integrated and referenced to quinine sulfate (quantum yield 0.51). A Hewlett-Packard Model 8452a spectrophotometer (Palo Alto, CA) was used for UV-VIS analysis. Lifetime measurements were obtained by fitting a multi-exponential model to time-correlated single photon counting decay data, using a weighted nonlinear least-square method (see, Grinvald *et al.*, *Anal. Biochem.*, **59**, 583-598, (1974)). Goodness of fit was assessed with the χ^2 function. (see, Badea *et al.*, *Methods in Enzymology*, **61**, 378-425, (1979)). For decay-associated spectra (DAS), time resolved data were obtained on samples by excitation at 330 nm and were observed every 10 nm over the emission band. The excitation pulse ("lamp") profile was obtained with a light-scattering suspension (Ludox) from Sigma. Further instrumentation details were as previously described. (see, Chen *et al. Biochemistry*, **30**, 5184-5195, (1991); Knutson, *et al. Biochemistry*, **21**, 4671-4679, (1982)). Deconvolution was routinely included in data analysis for the excitation pulse, which had an instrumental half-width of 800 ps (see, Chen *et al. Biochemistry*, **30**, 5184-5195, (1991)). A convolved multi-exponential model, $I'(t)$, describing the time course of fluorescence intensities (Eq. 1) was fit to the fluorescence decay data

$$I'(t) = \int L(t')I(t'-t)dt' \quad \text{Equation 1}$$

Where $L(t)$ is the lamp function (response of the instrument to the test laser pulse and

$$I(t) = \sum_{i=1}^n \alpha_i e^{(-t/\tau_i)} \quad \text{Equation 2}$$

Where $I(t)$ is the fluorescence intensity, α_i are the pre-exponentials, and τ_i are the lifetimes. Total fluorescence intensity, $I = \sum_{i=1}^n \alpha_i \tau_i$, and the percentage intensity from each component of the multi-exponential model is $\%I_i = \alpha_i \tau_i / I \times 100$. Mean lifetimes (intensity weighted lifetime, τ_m) and the species-concentration-weighted lifetime $\langle \tau \rangle$ are defined as:

$$\tau_m = \frac{\sum_{i=1}^n \alpha_i \tau_i^2}{\sum_{i=1}^n \alpha_i \tau_i} \quad \text{Equation 3}$$

And

$$\langle \tau \rangle = \frac{\sum_{i=1}^n \alpha_i \tau_i}{\sum_{i=1}^n \alpha_i} \quad \text{Equation 4}$$

Assessment of the degree of departure from mono-exponential decay was determined by comparing relative magnitudes of the pre-exponentials (α_i), the percentage contribution to the intensity (I_i) of each component, and the difference between τ_m and $\langle \tau \rangle$. When a decay is made up of components differing greatly in lifetime, τ_m will be much longer than $\langle \tau \rangle$.

For complex systems, DAS are the emission spectra that belong to each lifetime obtained from fluorescence decay surfaces (see, Knutson *et al.*, *Biochemistry*, **21**, 4671-4679, (1982)). DAS were used to dissect the heterogeneity of the emission but they do not, in themselves, specify its origin. Lifetimes were obtained by global analysis of the entire surface. Viewing DAS as plots of the pre-exponential constants (α_i) at each wavelength, these spectra can be normalized to provide the intensity contribution from each component. We evaluated the relationship between quantum yield and lifetimes, to better understand the mechanisms of quenching involved when pteridines are inserted into an oligonucleotide. For quenching events occurring during the excited state (pure dynamic quenching) where quenching competes with fluorescence,

$$Q = \frac{\tau}{\tau_n} \quad \text{Equation 5}$$

Where Q is the quantum yield, τ is the measured lifetime, and τ_n is the natural, or radiate, lifetime (*i.e.*, one that would be observed for $Q=1$). Deviations from Equation 5 such that $\tau/Q > \tau_n$ signify static (or quasi-static) quenching (*see*, Werner *et al.* 5 *Photochem. Photobiol.*, **29**, 905-914, (1979)), which in turn is usually due to ground-state formation of non-fluorescent complexes. Therefore, Equation 5 is an indicator of whether a quenching mechanism operates predominantly in the excited or ground state. For application to a heterogeneous solution, Eq. 5 may be modified to use a mean lifetime that is the sum of the contributions from each component, namely $\langle \tau \rangle$ (*see*, 10 Werner *et al.*, *Photochem. Photobiol.*, **29**, 905-914, (1979)).

RESULTS

Table 1

Fluorescence of DMAP and 6MAP monomers

	$Ex_{(max)}$	$Em_{(max)}$	Q_{rel}	$\tau(ns)$
6MAP compound 11	320	430	0.39	3.8
DMAP compound 12	310	430	0.48	4.8

Table 1: Fluorescence properties of the two adenosine analogues, 6MAP (compound 11) and DMAP (compound 12). Samples were measured in 10 mM Tris pH 7.5 at room temperature. Abbreviations: Ex_{max} , excitation maximum; Em_{max} , emission maximum; Q_{rel} , relative quantum yield; τ , lifetime in nanoseconds. 20

Relative quantum yields, excitation and emission maxima and lifetimes of the monomer form of 6MAP (compound 11) and DMAP (compound 12) are presented in Table 1. Each probe was found to be stable in ambient light and at room temperature for > 24 hours. Both of these compounds exhibited mono-exponential decay patterns in the 25 decay associated spectra.

Table 2

Relative quantum yields of oligonucleotides containing 6MAP or DMAP

SEQ ID:	Sequence (5' → 3')	w/6MAP	Q _{rel}
1	gtg tgg Faa atc tct agc agt	PTR21	0.010
2	gtg tgg aaa Ftc tct agc agt	PTR22	0.020
3	gtg tgg aaa atc tct Fgc agt	PTR23	0.018
4	act gct Fga gat ttt cca cac	PTR24	
5	act gct agF gat ttt cca cac	PTR25	0.011
6	act gct agc cFt ttt cca cac	PTR28	0.041
7	att cca caa Fgc cgt gtc a	HP21	0.010
8	aga ggt gtc cFc ctg tgg aga	HP22	<0.01
9	aga ggt gta cFa gtg tgg aga	HP23	0.012
10	aga ggt gta aFa atg tgg aga	HP24	<0.01
	Sequence (5' → 3')	w/DMAP	Q _{rel}
11	gtg tgg Faa atc tct agc agt	PTR31	0.023
12	gtg tgg aaa Ftc tct agc agt	PTR32	0.022
13	gtg tgg aaa atc tct Fgc agt	PTR33	0.01
14	act gct Fga gat ttt cca cac	PTR34	0.012
15	act gct agF gat ttt cca cac	PTR35	0.017
16	act gct aga gFt ttt cca cac	PTR36	0.019
17	act gct aga gat ttt ccF cac	PTR37	0.11
18	act gct agc cFt ttt cca cac	PTR38	0.11
19	att cca caa Fgc cgt gtc a	HP31	0.02
20	aga ggt gtc cFc ctg tgg aga	HP32	<0.01
21	aga ggt gta cFa gtg tgg aga	HP33	0.02
22	aga ggt gta aFa atg tgg aga	HP34	<0.01

5 Table 2: Relative quantum yields of oligo's listed in Table 2 were measured at room temperature in 10 mM Tris pH 7.5. The optical density of each was determined (at the excitation wavelength, 350 nm) and then each was scanned from 360 to 550 nm. The integral of the area under the emission curve was then used to calculate the relative quantum yield using quinine sulfate (Q= 0.51) as the standard.

10 Table 2: exhibits the relative quantum yields of a series of oligonucleotides containing either of the two probes. In each row, F denotes the position

of the probe within the sequence. Oligonucleotides containing 6MAP are numbered from 21 to 28 and those containing DMAP are numbered from 31 to 38. The sequences of the two series are identical. For example, PTR21 and PTR31 are identical in sequence, differing only in the identity of the probe (6MAP for PTR21 and DMAP for PTR31).

5

Table 3*Melting temperatures of Oligonucleotides containing 6MAP or DMAP*

Oligo w/6MAP	T _m	Oligo w/DMAP	T _m
PTR21	56.1°	PTR31	52.4°
PTR22	53.5°	PTR32	51.8°
PTR23	54.8°	PTR33	53.8°
PTR24	57.2°	PTR34	54.6°
PTR25	55.3°	PTR35	57.0°
		PTR36	51.6°
		PTR37	51.0°
		Control	57.8°

Table 3: Melting temperatures in degrees Centigrade. Samples were prepared from purified oligonucleotides diluted in 10 mM Tris pH 7.5 with 10 mM NaCl. Equimolar amounts of each strand were combined and heated to 85°C for 2 minutes and allowed to cool to room temperature. Melting temperatures of double-stranded oligonucleotides were measured by monitoring absorbance hyperchromicity at 260 nm. Melting temperatures of oligonucleotides containing either 6MAP or DMAP are shown in Table 3.

15

Each probe was tested for its ability to pair with bases other than thymidine. T_m's of the pairing with other bases are listed in Table 4.

Table 4*T_m's of the Pairing with other Bases*

	Paired w/T	Paired w/A	Paired w/C	Paired w/G
PTR22 (6MAP)	53.5°	43.8°	42.8°	45.2°
PTR32 (DMAP)	51.8°	45.2°	45.6°	46.8°

20

Table 4: Melting temperatures in degrees Centigrade for oligonucleotides containing 6MAP or DMAP paired with complementary strands with substitutions of T, A, C, or G as pairing partner for the probe. See Table 2 for sequences.

Table 5

5

Fluorescence of 6MAP or DMAP containing oligonucleotides

	Q_{rel}	τ_i (ns)	α_i	%I	τ_m (ns)	$\langle\tau\rangle$ (ns)
6MAP						
PTR21	0.01	$\tau_1=0.69$ $\tau_2=2.78$	$\alpha_1=0.30$ $\alpha_2=0.70$	30 70	2.58	2.15
PTR25	0.02	$\tau_1=2.93$				
PTR27		$\tau_1=0.29$ $\tau_2=1.54$ $\tau_3=4.73$	$\alpha_1=0.40$ $\alpha_2=0.5$ $\alpha_3=0.07$	40 53 7	2.26	1.26
PTR28	0.041	$\tau_1=0.17$ $\tau_2=1.15$ $\tau_3=2.56$	$\alpha_1=0.38$ $\alpha_2=0.40$ $\alpha_3=0.22$	38 40 22	1.82	1.09
HP21	0.01	$\tau_1=0.69$ $\tau_2=3.18$	$\alpha_1=0.41$ $\alpha_2=0.59$	41 59	2.85	2.16
HP22	>0.01	$\tau_1=0.61$ $\tau_2=3.12$	$\alpha_1=0.52$ $\alpha_2=0.48$	52 48	2.68	1.81
HP23	0.012	$\tau_1=0.48$ $\tau_2=2.58$	$\alpha_1=0.59$ $\alpha_2=0.41$	59 41	2.13	1.34
DMAP						
PTR32	0.02	$\tau_1=0.40$ $\tau_2=2.90$	$\alpha_1=0.39$ $\alpha_2=0.61$	39 61	2.7	1.92
PTR38	0.11	$\tau_1=0.77$ $\tau_2=2.80$	$\alpha_1=0.42$ $\alpha_2=0.58$	42 58	2.46	1.95
HP33	0.02	$\tau_1=0.28$ $\tau_2=2.52$	$\alpha_1=0.65$ $\alpha_2=0.35$	65 35	2.14	1.06

Table 5: Measurements were taken in 10 mM Tris buffer, pH 7.5 at room temperature. Abbreviations: Ex_{max} , excitation maximum; Em_{max} , emission maximum; Q_{rel} , relative quantum yield; τ_i , lifetime for each component of a multi-exponential model; α_i , pre-exponential for each component of a multi-exponential model; %I, percentage fluorescence intensity for each component of a multi-exponential model; $\langle\tau\rangle$, species-concentration-weighted lifetime; τ_m , intensity-weighted lifetime in nanoseconds.

P1 Nuclease Digestion of 6MAP or DMAP containing single strands

To determine that quenching within oligonucleotides is a function of incorporation into a strand, and not from degradation of the probe during DNA synthesis, a P1 nuclease digestion was performed on separate strands containing either 6MAP or DMAP. In each case the resulting increase in fluorescence intensity compared to the fluorescence intensity of the undigested equivalent formed a ratio that was equivalent to the ratio of the quantum yield for the single stranded oligonucleotide and the quantum yield for the monomer form.

The results as measured by fluorescence emission scan for P1 nuclease digestion of oligonucleotides containing either 6MAP or DMAP are shown in Figure 4 and Figure 5, respectively. Two separate strands for DMAP are compared to check the expected increase as compared to relative quantum yields for the oligonucleotide containing the fluorophore versus the relative quantum yield of the monomer form of the fluorophore. Real time data are displayed in Figure 6.


The monomer forms of 6MAP and DMAP display mono-exponential decay curves and in most cases, incorporation into an oligonucleotide results in an increase in complexity of the decay curves requiring two and sometimes three components to achieve an acceptable fit. This increase in complexity combined with the impact of severe decreases in the $\langle\tau\rangle$ (43 to 78%) is an indication of the degree of association of the pteridine analogues with the DNA. For both 6MAP and DMAP, quench associated with going into an oligonucleotide is >97% and the decrease in $\langle\tau\rangle$'s range from 43 to 71% for 6MAP and from 60% to 78% for DMAP. In those oligonucleotides experiencing a less severe decrease in $\langle\tau\rangle$, the quenching mechanism can be attributed mostly to static interactions. In general, it appears that the DMAP probe is more subject to dynamic quenching than 6MAP is since decreases in $\langle\tau\rangle$ are consistently greater.

These fluorophores, along with other fluorescent nucleoside analogues, exhibit unique properties that make them extremely valuable for measuring subtle events within DNA. They are highly fluorescent, stable and easy to use. Even though these compounds are significantly quenched when incorporated into an oligonucleotide, they
5 are still very detectable using a standard bench top fluorometer. Additionally, placing one strategically within a strand of DNA gives a picture of events occurring close to that site as reported directly by changes in fluorescence intensity, anisotropy, energy transfer, or lifetime measurements experienced by the probe.

All publications, patents and patent applications mentioned in this
10 specification are herein incorporated by reference into the specification in their entirety for all purposes. Although the invention has been described with reference to preferred embodiments and examples thereof, the scope of the present invention is not limited only to those described embodiments. As will be apparent to persons skilled in the art, modifications and adaptations to the above-described invention can be made without
15 departing from the spirit and scope of the invention, which is defined and circumscribed by the appended claims.

WHAT IS CLAIMED IS:

1. A compound of the formula:



wherein:


R^1 is a member selected from the group consisting of hydrogen and optionally substituted C_1 - C_6 -alkyl;

R^2 is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R^3 is optionally substituted C_1 - C_6 alkyl;

R^4 is a member selected from the group consisting of hydrogen and L;

L is of the formula



wherein:

R^5 is a member selected from the group consisting of hydrogen, hydroxyl, and substituted hydroxyl wherein the substituent is a protecting group;

R^6 is a member selected from the group consisting of hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support, a hydroxyalkyl, and a hydroxyalkyl covalently bound to a solid support; and

R^7 is a member selected from the group consisting of hydrogen, a phosphate, a triphosphate, and a protecting group;

with the proviso that R^1 and R^4 are not simultaneously L.

2. A compound in accordance with claim 1, wherein R^1 is hydrogen;

R^2 is a member selected from the group consisting of amino, mono-, and di-substituted amino wherein the substituents are members selected from the group consisting of benzoyl, isobutyryl, phthaloyl, di-n-butylaminomethylidene,

5 dimethylaminomethylidene, p-nitrophenylethoxycarbonyl and
6 dimethylaminomethylenamino;
7 R^4 is L;
8 R^5 is a member selected from the group consisting of hydrogen, hydroxyl,
9 hydroxyl substituted with a member selected from the group consisting of trityl,
10 monomethoxytrityl, dimethoxytrityl, tetrahydropyran-1-yl, 4-methoxytetrahydropyran-4-
11 yl, 1-(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl, t-butyldimethylsilyl, p-
12 nitrophenylethylsulfonyl, tetrahydropyranyl, 4-methoxytetrahydropyranyl, 2-nitrobenzyl,
13 9-phenylxanthen-9-yl and p-nitrophenylethyl;
14 R^6 is a member selected from the group consisting of consisting of
15 hydrogen, phosphoramidite, H-phosphonate, hemisuccinate, and hemisuccinate
16 covalently bound to a solid support; and
17 R^7 is a member selected from the group consisting of hydrogen, trityl,
18 monomethoxytrityl, dimethoxytrityl, phthaloyl, di-n-butylaminomethylene,
19 dimethylaminomethylidene and triphosphate.

1 3. A compound in accordance with claim 2, wherein R^2 is a member
2 selected from the group consisting of amino and an amino group mono-substituted by a
3 protecting group selected from the group consisting of di-n-butylaminomethylidene, p-
4 nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

5 R^5 is a member selected from the group consisting of hydrogen, hydroxyl
6 and hydroxyl substituted with a member selected from the group consisting of
7 dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-
8 nitrophenylethylsulfonyl;

9 R^6 is a member selected from the group consisting of hydrogen, β -
10 cyanoethyl-N-diisopropyl phosphoramidite and a hemisuccinate covalently bound to
11 controlled pore glass; and

12 R^7 is a member selected from the group consisting of dimethoxytrityl, di-
13 n-butylaminomethylene, and dimethylaminomethylidene.

1 4. A compound in accordance with claim 2, wherein R^2 is a member
2 selected from the group consisting of amino and an amino group mono-substituted by a
3 protecting group selected from the group consisting of di-n-butylaminomethylidene, p-
4 nitrophenylethoxycarbonyl, and dimethylaminomethylenamino;

5 R⁵ is a member selected from the group consisting of hydrogen and
6 hydroxyl substituted with a member selected from the group consisting of
7 dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-
8 nitrophenylethyl;

9 R⁶ is a member selected from the group consisting of hydrogen and β -
10 cyanoethyl-N-diisopropyl phosphoramidite; and

11 R⁷ is a member selected from the group consisting of hydrogen and
12 dimethoxytrityl.

1 5. A compound in accordance with claim 2, wherein R² is a member
2 selected from the group consisting of amino and dimethylaminomethylenamino;

3 R³ is methyl;

4 R⁵ is hydrogen;

5 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

6 R⁷ is dimethoxytrityl.

1 6. A compound in accordance with claim 2, wherein R² is amino;

2 R³ is methyl;

3 R⁵ is hydrogen;

4 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

5 R⁷ is dimethoxytrityl.

1 7. A compound in accordance with claim 2, wherein R² is amino;

2 R³ is methyl;

3 R⁵ is hydrogen;

4 R⁶ is hydrogen; and

5 R⁷ is hydrogen.

1 8. A compound in accordance claim 2, wherein R² is

2 dimethylaminomethylenamino;

3 R³ is methyl;

4 R⁵ is hydrogen;

5 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

6 R⁷ is dimethoxytrityl.

1 9. A compound in accordance with claim 2, wherein R² is amino;

2 R^3 is methyl;
3 R^5 is hydrogen;
4 R^6 is β -cyanoethyl-N-diisopropyl phosphoramidite; and
5 R^7 is a triphosphate.

1 10. A compound in accordance with claim 1, wherein;
2 R^1 is optionally substituted C_1 - C_6 alkyl;
3 R^2 is a member selected from the group consisting of amino, mono-, and
4 di-substituted amino wherein the substituent is a member selected from the group
5 consisting of benzoyl, isobutyryl, phthaloyl, di-n-butylaminomethylidene,
6 dimethylaminomethylidene, p-nitrophenylethoxycarbonyl and
7 dimethylaminomethylenamino;
8 R^3 is optionally substituted C_1 - C_6 alkyl;
9 R^4 is L;
10 R^5 is a member selected from the group consisting of hydrogen, hydroxyl
11 and hydroxyl substituted with a member selected from the group consisting of trityl,
12 monomethoxytrityl, dimethoxytrityl, tetrahydropyran-1-yl, 4-methoxytetrahydropyran-4-
13 yl, 1-(2-chloro-4-methyl)phenyl-4-methoxypiperidin-4-yl, t-butyldimethylsilyl, p-
14 nitrophenylethylsulfonyl, tetrahydropyranyl, 4-methoxytetrahydropyranyl, 2-nitrobenzyl,
15 9-phenylxanthen-9-yl and p-nitrophenylethyl;
16 R^6 is a member selected from the group consisting of hydrogen, H-
17 phosphonate, phosphoramidite, hemisuccinate, and hemisuccinate covalently bound to a
18 solid support; and
19 R^7 is a member selected from the group consisting of hydrogen, trityl,
20 monomethoxytrityl, dimethoxytrityl, phthaloyl, di-n-butylaminomethylene, and
21 dimethylaminomethylidene.

1 11. A compound in accordance with claim 10 wherein R^1 is methyl;
2 R^2 is a member selected from the group consisting of amino and an amino
3 group mono-substituted by a protecting group selected from the group consisting of di-n-
4 butylaminomethylidene, p-nitrophenylethoxycarbonyl, and
5 dimethylaminomethylenamino;
6 R^3 is methyl;

7 R⁵ is a member selected from the group consisting of hydrogen, hydroxyl
8 and hydroxyl substituted with a member selected from the group consisting of
9 dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-
10 nitrophenylethylsulfonfyl;

11 R⁶ is a member selected from the group consisting of hydrogen, β -
12 cyanoethyl-N-diisopropyl phosphoramidite and a hemisuccinate covalently bound to
13 controlled pore glass; and

14 R⁷ is a member selected from the group consisting of dimethoxytrityl, di-
15 n-butylaminomethylene, and dimethylaminomethylidene.

1 12. A compound in accordance claim 10, wherein R¹ is methyl; R² is a
2 member selected from the group consisting of amino and an amino group mono-
3 substituted by a protecting group selected from the group consisting of di-n-
4 butylaminomethylidene, p-nitrophenylethoxycarbonyl, and
5 dimethylaminomethylenamino;

6 R⁵ is a member selected from the group consisting of hydrogen and
7 hydroxyl substituted with a member selected from the group consisting of
8 dimethoxytrityl, tetrahydropyran-1-yl, t-butyldimethylsilyl, 2-nitrobenzyl, and p-
9 nitrophenylethylsulfonfyl;

10 R⁶ is a member selected from the group consisting of consisting of
11 hydrogen and β -cyanoethyl-N-diisopropyl phosphoramidite; and

12 R⁷ is a member selected from the group consisting of hydrogen and
13 dimethoxytrityl.

1 13. A compound in accordance with claim 10, wherein R¹ is methyl;
2 R² is a member selected from the group consisting of amino and
3 dimethylaminomethylenamino;

4 R³ is methyl;

5 R⁵ is hydrogen;

6 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and

7 R⁷ is dimethoxytrityl.

1 14. A compound in accordance with claim 10, wherein R¹ is methyl;
2 R² is amino;

3 R³ is methyl;


4 R⁵ is hydrogen;
 5 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and
 6 R⁷ is dimethoxytrityl.

1 15. A compound in accordance with claim 10, wherein R¹ is methyl;
 2 R² is amino;
 3 R³ is methyl;
 4 R⁵ is hydrogen;
 5 R⁶ is hydrogen; and
 6 R⁷ is hydrogen.

1 16. A compound in accordance with claim 10, wherein R¹ is methyl; is
 2 dimethylaminomethylenamino;
 3 R³ is methyl;
 4 R⁵ is hydrogen;
 5 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and
 6 R⁷ is dimethoxytrityl.


1 17. A compound in accordance with claim 10, wherein R¹ is methyl;
 2 R² is amino;
 3 R³ is methyl;
 4 R⁵ is hydrogen;
 5 R⁶ is β -cyanoethyl-N-diisopropyl phosphoramidite; and
 6 R⁷ is a triphosphate.

1 18. An oligonucleotide comprising one or more nucleotide monomers,
 2 said monomers having the formula



3
 4 wherein:
 5 R¹ is a member selected from the group consisting of hydrogen and
 6 optionally substituted C₁-C₆-alkyl;

- 7 R^2 is a member selected from the group consisting of amino and mono- or
 8 di-substituted amino wherein the substituent is a protecting group;
 9 R^3 is optional substituted C_1 - C_6 alkyl;
 10 R^4 is L;
 11 L is of the formula



- 12
 13 wherein:
 14 R^5 is a member selected from the group consisting of hydrogen and
 15 hydroxyl;
 16 R^6 is a member selected from the group consisting of hydrogen, a
 17 phosphate, a phosphate covalently attached to a nucleotide, a phosphate covalently
 18 attached to a nucleoside; a hemisuccinate covalently bound to a solid support, a
 19 dicyclohexylcarbodiimide covalently bound to a solid support, and a hydroxyalkyl
 20 covalently bound to a solid support; and
 21 R^7 is a member selected from the group consisting of hydrogen, a
 22 phosphate, a phosphate covalently attached to a nucleotide and a phosphate covalently
 23 attached to a nucleoside;
 24 with the proviso that R^1 and R^4 are not simultaneously L.

- 1 19. An oligonucleotide in accordance with claim 18, wherein:
 2 R^1 is hydrogen,
 3 R^2 is amino;
 4 R^3 is methyl;
 5 R^5 is hydrogen and hydroxyl;
 6 R^6 is hydrogen; and
 7 R^7 is a phosphate.

- 1 20. An oligonucleotide in accordance with claim 19, wherein:
 2 R^5 is hydrogen.

- 1 21. An oligonucleotide in accordance with claim 19 wherein:

2 R⁵ is hydroxyl.

1 22 An oligonucleotide in accordance with claim 18, wherein:

2 R¹ is optionally substituted C₁-C₆-alkyl;

3 R² is amino;

4 R³ is methyl;

5 R⁵ is hydrogen and hydroxyl;

6 R⁶ is hydrogen; and

7 R⁷ is a phosphate.

1 23. An oligonucleotide in accordance with claim 22, wherein

2 R¹ is methyl and

3 R⁵ is hydrogen.

1 24. An oligonucleotide in accordance with claim 22, wherein

2 R¹ is methyl and

3 R⁵ is hydroxyl.

1 25. An oligonucleotide in accordance with claim 18, wherein said

2 nucleotide monomers are at the 3' end of said oligonucleotide.

1 26. An oligonucleotide in accordance with claim 18, wherein said

2 nucleotide monomers are at the 5' end of said oligonucleotide.

1 27. An oligonucleotide in accordance with claim 18, wherein said

2 nucleotide monomers are surrounded by 1 to 10 pyrimidine monomers.

1 28. An oligonucleotide in accordance with claim 18, wherein said

2 oligonucleotide is a member selected from the group consisting of SEQ ID:1, SEQ ID:2,

3 SEQ ID:3, SEQ ID:4, SEQ ID:5, SEQ ID:6, SEQ ID:7, SEQ ID:8, SEQ ID:9, SEQ ID:10,

4 SEQ ID:11, SEQ ID:12, SEQ ID:13, SEQ ID:14, SEQ ID:15, SEQ ID:16, SEQ ID:17,


5 SEQ ID:18, SEQ ID:19, SEQ ID:20, SEQ ID:21 and SEQ ID:22.

1 29. A method of detecting the presence, absence, or quantity of a target

2 nucleic acid, said method comprising the steps of:

3 a) contacting said target nucleic acid with a nucleic acid probe wherein said

4 nucleic acid probe comprises compound of the formula:



wherein:


R^1 is a member selected from the group consisting of hydrogen and optionally substituted C_1 - C_6 -alkyl;

R^2 is a member selected from the group consisting of amino and mono- or di-substituted amino wherein the substituent is a protecting group;

R^3 is optionally substituted C_1 - C_6 alkyl;

R^4 is L;

L is of the formula



wherein:

R^5 is a member selected from the group consisting of hydrogen and hydroxyl;

R^6 is a member selected from the group consisting of hydrogen, phosphoramidite, an H-phosphonate, a methyl phosphonate, a phosphorothioate, a phosphotriester, a hemisuccinate, a hemisuccinate covalently bound to a solid support, a dicyclohexylcarbodiimide, and a dicyclohexylcarbodiimide covalently bound to a solid support; and

R^7 is phosphate;

with the proviso that R^1 and R^4 are not simultaneously L; located in said probe such that, when said probe hybridizes to said target nucleic acid said compound is in a loop that does not participate in complementary base pairing with a nucleotide of said target nucleic acid; and

b) detecting the fluorescence produced by said fluorescent nucleotide when said probe forms a hybrid duplex with said target nucleic acid.

30. A method of claim 29, wherein said loop ranges in length from about 1 to about 100 nucleotides when said probe hybridizes to said target nucleic acid

1 31. A method of claim 29, wherein said loop is an insertion in said
2 nucleic acid probe which is otherwise complementary to said target nucleic acid or to a
3 contiguous subsequence of said target nucleic acid.

1 32. A method of claim 31, wherein said insertion is three nucleotides in
2 length and comprises two nucleotides each adjacent to said compound.

1 33. A method of claim 32, wherein at least one nucleotide adjacent to
2 said compound is a purine.

1 34. A method of claim 33, wherein at least one nucleotide adjacent to
2 said compound is an adenosine.

1 35. A method of claim 32, wherein at least one nucleotide adjacent to
2 said compound is a pyrimidine.

1 36. A method of claim 35, wherein at least one nucleotide adjacent to
2 said compound is a cytosine.

1 37. A method of claim 34, wherein said compound is bordered by at
2 least two adjacent purines in both the 5' and 3' direction.

1 38. A method of claim 37, wherein said adjacent purines are adenosine.

1 39. A method of claim 31, wherein said insertion is said compound.

1 40. A method of claim 31, wherein said insertion is self-
2 complementary and forms a hairpin wherein said compound is present in the loop of said
3 hairpin and does not participate in complementary base pairing.

1 41. A method of claim 29, wherein the nucleotides comprising said
2 loop are selected such that they are not complementary to the corresponding nucleotides
3 of the target nucleic acid when said probe is hybridized to said target nucleic acid and
4 wherein said probe is complementary to at least two non-contiguous subsequences of said
5 target nucleic acid.

1 42. A method of claim 29, wherein said fluorescent nucleotide is
2 present in a terminal subsequence of said nucleic acid probe wherein said terminal

3 subsequence does not hybridize to said target nucleic acid when the remainder of said
4 nucleic acid probe hybridizes to said target nucleic acid.

1 43. A method of claim **42**, wherein said terminal subsequence forms a
2 terminal hairpin by hybridization with a second subsequence of said probe such that said
3 fluorescent nucleotide is present in a loop of said hairpin and does not participate in
4 complementary base pairing.

1 44. A method of claim **29**, wherein said detecting comprises detecting
2 an increase in fluorescence of said fluorescent nucleotide when said probe forms a hybrid
3 duplex with said target nucleic acid.

1 45. A kit for the detection of nucleic acid-nucleic acid interactions
2 comprising a container, said container containing a compound in accordance with claim
3 **1**, and instructions for use.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: PTERIDINE NUCLEOTIDE ANALOGS, the specification of which was filed on 07 September 1999 as PCT International Application No. PCT/US99/20541, and which received U.S. Serial Number 09/786,666.

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Country	Application No.	Date of Filing	Priority Claimed Under 35 USC 119
			Yes <input type="checkbox"/> No <input type="checkbox"/>
			Yes <input type="checkbox"/> No <input type="checkbox"/>

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

Application No.	Filing Date
60/099,487	08 September 1998

I claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Date of Filing	Status
		<input type="checkbox"/> Patented <input type="checkbox"/> Pending <input type="checkbox"/> Abandoned
		<input type="checkbox"/> Patented <input type="checkbox"/> Pending <input type="checkbox"/> Abandoned
		<input type="checkbox"/> Patented <input type="checkbox"/> Pending <input type="checkbox"/> Abandoned

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) of the Government of the United States of America as represented by the Secretary of the Department of Health and Human Services, Office of Technology Transfer, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.


Robert Benson, Reg. No. 33,612
Steven M. Ferguson, Reg. No. 38,448
James C. Haight, Reg. No. 25,588
John P. Kim, Reg. No. 38,514

Susan S. Rucker, Reg. No. 35,762
David R. Sadowski, Reg. No. 32,808
Jack Spiegel, Reg. No. 34,477

Send Correspondence to Appointed Associate Attorney or Agent Address: Eugenia Garrett-Wackowski TOWNSEND and TOWNSEND and CREW LLP Two Embarcadero Center, 8th Floor San Francisco, CA 94111-3834	Direct Telephone Calls and Facsimiles to Appointed Associate Attorney or Agent: Name: Eugenia Garrett-Wackowski Reg. No.: 37,330 Telephone: 415/576-0200 Fax: 415/576-0300
---	--

Full Name of Inventor 1	Last Name Hawkins	First Name Mary	Middle Name or Initial E.	
Residence & Citizenship	City Potomac	State/Foreign Country Maryland	Country of Citizenship US	
Post Office Address	Post Office Address 5 Eldwick Court	City Potomac	State/Country Maryland	Zip Code 20854
Full Name of Inventor 2	Last Name Pfleiderer	First Name Wolfgang	Middle Name or Initial	
Residence & Citizenship	City Konstanz	State/Foreign Country Germany	Country of Citizenship DE	
Post Office Address	Post Office Address Lindauerstrasse 47, 78464	City Konstanz	State/Country Germany	Zip Code
Full Name of Inventor 3	Last Name Balis	First Name Frank	Middle Name or Initial	
Residence & Citizenship	City Bethesda	State/Foreign Country Maryland	Country of Citizenship US	
Post Office Address	Post Office Address 6310 Rockhurst Road	City Bethesda	State/Country Maryland	Zip Code 20817


I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18 of the United States Code, Section 1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2	Signature of Inventor 3  5/10/01
Date	Date	Date

Send Correspondence to Appointed Associate Attorney or Agent Address: Eugenia Garrett-Wackowski TOWNSEND and TOWNSEND and CREW LLP Two Embarcadero Center, 8th Floor San Francisco, CA 94111-3834	Direct Telephone Calls and Facsimiles to Appointed Associate Attorney or Agent: Name: Eugenia Garrett-Wackowski Reg. No.: 37,330 Telephone: 415/576-0200 Fax: 415/576-0300
---	--

Full Name of Inventor 1	Last Name Hawkins	First Name Mary	Middle Name or Initial E.	
Residence & Citizenship	City Potomac	State/Foreign Country Maryland	Country of Citizenship US	
Post Office Address	Post Office Address 5 Eldwick Court	City Potomac	State/Country Maryland	Zip Code 20854
Full Name of Inventor 2	Last Name Pfleiderer	First Name Wolfgang	Middle Name or Initial	
Residence & Citizenship	City Konstanz	State/Foreign Country Germany	Country of Citizenship DE	
Post Office Address	Post Office Address Lindauerstrasse 47, 78464	City Konstanz	State/Country Germany	Zip Code
Full Name of Inventor 3	Last Name Balis	First Name Frank	Middle Name or Initial	
Residence & Citizenship	City Bethesda	State/Foreign Country Maryland	Country of Citizenship US	
Post Office Address	Post Office Address 6310 Rockhurst Road	City Bethesda	State/Country Maryland	Zip Code 20817


I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18 of the United States Code, Section 1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2 	Signature of Inventor 3
Date	Date <u>5-14-2001</u>	Date

Send Correspondence to Appointed Associate Attorney or Agent Address: <u>Eugenia Garrett-Wackowski</u> <u>TOWNSEND and TOWNSEND and CREW LLP</u> <u>Two Embarcadero Center, 8th Floor</u> <u>San Francisco, CA 94111-3834</u>	Direct Telephone Calls and Facsimiles to Appointed Associate Attorney or Agent: Name: Eugenia Garrett-Wackowski Reg. No.: 37,330 Telephone: 415/576-0200 Fax: 415/576-0300
---	--

Full Name of Inventor 1	Last Name <u>Hawkins</u>	First Name <u>Mary</u>	Middle Name or Initial <u>E.</u>	
Residence & Citizenship	City <u>Potomac</u>	State/Foreign Country <u>Maryland</u> <u>MD</u>	Country of Citizenship <u>US</u>	
Post Office Address	Post Office Address <u>5 Eldwick Court</u>	City <u>Potomac</u>	State/Country <u>Maryland</u>	Zip Code <u>20854</u>
Full Name of Inventor 2	Last Name <u>Pfleiderer</u>	First Name <u>Wolfgang</u>	Middle Name or Initial	
Residence & Citizenship	City <u>Konstanz</u>	State/Foreign Country <u>Germany</u> <u>DEX</u>	Country of Citizenship <u>DE</u>	
Post Office Address	Post Office Address <u>Lindauerstrasse 47, 78464</u>	City <u>Konstanz</u>	State/Country <u>Germany</u>	Zip Code
Full Name of Inventor 3	Last Name <u>Balis</u>	First Name <u>Frank</u>	Middle Name or Initial	
Residence & Citizenship	City <u>Bethesda</u>	State/Foreign Country <u>Maryland</u> <u>MD</u>	Country of Citizenship <u>US</u>	
Post Office Address	Post Office Address <u>6310 Rockhurst Road</u>	City <u>Bethesda</u>	State/Country <u>Maryland</u>	Zip Code <u>20817</u>

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18 of the United States Code, Section 1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1 	Signature of Inventor 2	Signature of Inventor 3
Date <u>5-10-2001</u>	Date	Date